

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/62, C07K 16/10, C12N 15/13, 15/70, 15/85, 1/21, 5/10, A61K 39/42, G01N 33/577, 33/569, A61K 47/48, 38/20, 31/70 // (A61K 39/42, 38:21) (A61K 39/42, 31:70)	A1	(11) International Publication Number: WO 97/40176 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/EP97/01977 (22) International Filing Date: 18 April 1997 (18.04.97) (30) Priority Data: 08/635,109 19 April 1996 (19.04.96) US 08/844,215 17 April 1997 (17.04.97) US (71)(72) Applicants and Inventors: PERSSON, Mats, Axel, Atterdag [SE/SE]; Karolinska Hospital, Dept. of Medicine M3:00, S-171 76 Stockholm (SE). ALLANDER, Tobias Erik [SE/SE]; Karolinska Hospital, Dept. of Medicine M3:00, S-171 76 Stockholm (SE). (74) Agents: GOLDIN, Douglas, M.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5LX (GB) et al.		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HUMAN MONOCLONAL ANTIBODIES SPECIFIC FOR HEPATITIS C VIRUS (HCV) E2 ANTIGEN (57) Abstract <p>The present invention relates to compositions derived from immunoglobulin molecules specific for the hepatitis C virus (HCV). More particularly, the invention is related to molecules which are capable of specifically binding with HCV E2 antigen. The molecules are useful in specific binding assays, affinity purification schemes and pharmaceutical compositions for the prevention and treatment of HCV infection in mammalian subjects. The invention thus relates to novel human monoclonal antibodies specific for HCV E2 antigen, fragments of such monoclonal antibodies, polypeptides having structure and function substantially homologous to antigen-binding sites obtained from such monoclonal antibodies, nucleic acid molecules encoding those polypeptides, and expression vectors comprising the nucleic acid molecules.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5 HUMAN MONOCLONAL ANTIBODIES SPECIFIC FOR
 HEPATITIS C VIRUS (HCV) E2 ANTIGEN

Technical Field

10 The present invention relates to
 compositions derived from immunoglobulin molecules
 specific for the hepatitis C virus (HCV). More
 particularly, the invention is related to recombinant
 human monoclonal antibodies which are capable of
15 specifically binding with HCV E2 antigen.

Background

 Hepatitis C virus (HCV) infection occurs
 throughout the world and is the major cause of
20 transfusion-associated hepatitis. There are an
 estimated 150,000 new cases of HCV infection each year
 in the United States. The seroprevalence of anti-HCV
 antibodies in blood donors from around the world has
 been shown to vary between 0.02 and 1.23%, with rates
25 in some countries as high as about 19%. In addition
 to being the predominate cause of transfusion-induced
 hepatitis, HCV is also a common cause of hepatitis in
 individuals exposed to blood or blood products. Thus,
 recipients of blood or blood products, intravenous
30 drug users, renal dialysis patients and needle-stick
 victims represent high-risk groups for HCV infection.
 Alter et al. (1993) *Infect Agents Dis* 2:155-166.
 Further, heterosexual transmission of HCV across the
 urogenital tract, and mother-to-baby transmission, has
35 been well documented. Ohto et al. (1994) *N Engl J Med*
 330:744-750. Other risk factors associated with HCV
 infection include familial or household contact with

an HCV-infected individual and health-care employment with occupational exposure to blood and hemodialysis. Alter et al. (1990) *JAMA* 264:2231-2235. Chronic hepatitis develops in approximately 62% of infections.

5 Alter et al. (1992) *N Engl J Med* 327:1899-1905.

Most of the serious liver disease associated with HCV results from the high propensity of the agent to cause chronic, persistent infection. Cirrhosis occurs in approximately 20% of chronic cases, of which
10 20 to 25% will result in liver failure. Another serious sequela associated with HCV infection is primary hepatocellular carcinoma.

The viral genomic sequence of HCV is known, as are methods for obtaining the sequence. See, e.g.,
15 International Publication Nos. WO 89/04669; WO 90/11089; and WO 90/14436. HCV has a 9.5 kb positive-sense, single-stranded RNA genome and is a member of the Flaviridae family of viruses. Currently, there are 6 distinct, but related genotypes of HCV which
20 have been identified based on phylogenetic analyses (Simmonds et al., *J. Gen. Virol.* (1993) 74:2391-2399). The virus encodes a single polypeptide having more than 3000 amino acid residues (Choo et al. (1989) *Science* 244:359-362; Choo et al. (1991) *Proc. Natl.*
25 *Acad. Sci. USA* 88:2451-2455; Han et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:1711-1715). The polypeptide is processed co- and post-translationally into both structural and non-structural (NS) proteins.

In particular, there are three putative HCV
30 structural proteins, consisting of the N-terminal nucleocapsid protein (termed "core") and two envelope glycoproteins, "E1" (also known as E) and "E2" (also known as E2/NS1). (See, Houghton et al. (1991) *Hepatology* 14:381-388, for a discussion of HCV
35 proteins, including E1 and E2.) E1 is detected as a 32-35 kDa species and is converted into a single endo H-sensitive band of approximately 18 Kda. By

contrast, E2 displays a complex pattern upon immunoprecipitation consistent with the generation of multiple species (Grakoui et al. (1993) *J. Virol.* 67:1385-1395; Tomei et al. (1993) *J. Virol.* 67:4017-4026).

5 The HCV envelope glycoproteins E1 and E2 form a stable complex that is coimmunoprecipitable (Grakoui et al. (1993) *J. Virol.* 67:1385-1395; Lanford et al. (1993) *Virology* 197:225-235; Ralston et al. (1993) *J. Virol.* 67:6753-6761).

10 The only currently available treatment for chronic hepatitis C infection consists of α -interferon (α -IFN) therapy. However, long-term response to interferon therapy only occurs in 10% to 30% of treated individuals, and there is evidence that the
15 different HCV strains vary greatly in their responsiveness to interferon therapy, with the type 1 viruses being the most refractive. Furthermore, flu-like side effects are commonly encountered with interferon therapy (occurring in approximately 60% to
20 80% of treated individuals), as well as other less common side effects such as nausea, depression, fatigue and thrombocytopenia. Interferon therapy is also not indicated for immunocompromised individuals. Accordingly, there exists a need for more effective
25 therapeutic approaches in the treatment of chronic HCV infection. In this regard, some effect has been seen using ribivirin, or combination therapies with ursodiol and α -IFN.

 In particular, the HCV E1 and E2 proteins
30 are of considerable interest because recombinant vaccines based on those molecules have been shown to be protective against experimental challenge with HCV in primate studies. (Choo et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:1294-1298). Hyperimmune globulin
35 compositions of anti-HCV antibody molecules obtained from donor samples have been described for the treatment of HCV in infected individuals, and in the

prevention of HCV infection in high-risk groups.
European Patent Application Publication No. 447,984,
published 25 September 1991. Since these compositions
are made from donor blood products, an inherent risk
5 is associated with their use due to the possible
presence of infectious agents such as the Human
Immunodeficiency Virus (HIV) and HCV. Accordingly,
hyperimmune globulin preparations must be carefully
screened, and all infectious agents inactivated prior
10 to administration to human subjects.

It is known that the immune response to HCV
in normal individuals includes both humoral and cell
mediated components. Koziel et al. (1993) *J Virol*
67:7522-7532, Alter et al. (1989) *N Engl J Med*
15 321:1494-1500. Further, several reports have
indicated that antibodies elicited to HCV may
neutralize the infectivity of the virus. Shimizu et
al. (1994) *J Virol* 68:1494-1500, Farci et al. (1994)
Proc Natl Acad Sci USA 91:7792-7796. Such results
20 provide hope that an effective antibody-based therapy
can be developed. In this regard, the administration
of a highly-reactive, neutralizing anti-HCV antibody
preparation to an individual who is at risk of
infection, or who has been recently exposed to the
25 agent will provide immediate passive immunity to the
individual. Such passive immunizations would likewise
be expected to be successful in both normal and
immunocompromised subjects. Preferably, the
neutralizing antibodies would be broadly cross-
30 reactive against different HCV strains, and would be
monoclonal in order to control the effects of the use
of the antibodies *in vivo*.

For a number of practical and economic
reasons, murine monoclonal antibodies have been
35 generally used in research and medicine. Murine
antibodies can be raised against a wide variety of
molecules, such as HCV antigens, and fused with a

myeloma cell to yield hybridomas which can be grown in culture to produce monoclonal antibodies toward HCV antigens. Kohler et al. (1975) *Nature* 256:495-497. Although such monoclonal antibodies may have antigen
5 binding specificities of significant therapeutic value, the use of such murine antibodies in the treatment of human disease has been limited since those molecules are immunogenic to the human immune system. Thus, murine monoclonals have been most
10 commonly used in immunodiagnostics. In this regard, murine monoclonal antibodies to putative HCV E2 envelope polypeptides have been described for use in the detection of HCV in biological samples. U.S. Patent No. 5,308,750 to Mehta et al.

15 Accordingly, there remains a need in the art to provide human monoclonal antibodies toward HCV E2 antigen, wherein the monoclonals are broadly cross-reactive with heterologous HCV isolates.

20 Summary of the Invention

The present invention is based on the discovery of human monoclonal antibody molecules which exhibit immunological binding affinity for HCV E2 polypeptide antigen, and which are cross-reactive
25 against different HCV strains. The monoclonal antibody molecules were obtained from a combinatorial library that was constructed from a nonimmunized HCV-infected source. The present molecules generally comprise a human antibody Fab molecule that exhibits
30 immunological binding affinity for HCV E2 antigen.

Accordingly, in one embodiment, the invention is directed to a recombinant human monoclonal antibody that exhibits immunological binding affinity for HCV E2 antigen, wherein the
35 monoclonal antibody includes amino acid sequences that are homologous to the binding portion of a human antibody Fab molecule obtained from a combinatorial

antibody library. The recombinant monoclonal antibody molecule can be in the form of a substantially whole immunoglobulin molecule, or can be in the form of a soluble Fab molecule, an Fv fragment, or an sFv molecule, wherein each molecule at least contains amino acid sequences that are homologous to the binding portion of a human antibody Fab molecule.

In another embodiment, the invention is directed to an isolated nucleic acid molecule which contains a polynucleotide coding sequence for a polypeptide that is homologous to the binding portion of a heavy or light chain variable region (V_H or V_L) of a human Fab molecule which exhibits immunological binding affinity for HCV E2 antigen. In a related embodiment, the invention is directed to an isolated nucleic acid molecule which contains polynucleotide coding sequences for a first polypeptide and polynucleotide coding sequences for a second polypeptide, wherein the first polypeptide is homologous to the binding portion of a heavy chain variable region (V_H) of a human Fab molecule which exhibits immunological binding affinity for HCV E2 antigen, and the second polypeptide is homologous to the binding portion of a light chain variable region (V_L) of a human Fab molecule which exhibits immunological binding affinity for the HCV E2 antigen.

In other embodiments, the invention pertains to expression vectors comprising the nucleic acid molecules above operably linked to control sequences that direct the transcription of the polynucleotide coding sequences when the vector is present in a host cell or under suitable conditions for the transcription and translation of the polynucleotide coding sequences. Yet further embodiments of the invention pertain to host cells transformed with the vectors of the invention, and methods for producing

recombinant polypeptides using the transformed host cells.

In another embodiment, the invention is directed to vaccine compositions comprising the recombinant monoclonal antibody molecules of the invention. Still further embodiments relate to methods of using the vaccine compositions, wherein the vaccines are used to provide an antibody titer to HCV in a mammalian subject, and/or used to provide passive immunity against HCV infection in a vaccinated subject. In related embodiments, the vaccine compositions are used in combination with known anti-HCV therapeutics.

In still further embodiments, the recombinant monoclonal antibody molecules of the invention are used to provide binding complexes which are labeled with a detectable moiety. The labeled binding complexes are used in related embodiments of the invention, such as in specific binding assay methods, for detecting the presence of HCV particles in samples suspected of containing HCV and in specific binding assays for monitoring the progress of anti-HCV treatment of HCV-infected subjects.

These and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

Brief Description of the Figures

Figures 1A-1G depict the deduced γ 1 heavy chain amino acid sequences of the Fab molecule clones 1:5 (SEQ ID NO:), 1:7 (SEQ ID NO:), 1:11 (SEQ ID NO:), L3 (SEQ ID NO:), L1 (SEQ ID NO:), A8 (SEQ ID NO:), and A12 (SEQ ID NO:), respectively. The CDR regions of each heavy chain have been identified in the Figures as "CDR1," "CDR2" and "CDR3."

Figures 2A-2G depict the deduced κ light chain amino acid sequences of the Fab molecule clones

1:5 (SEQ ID NO:), 1:7 (SEQ ID NO:), 1:11 (SEQ ID NO:), L3 (SEQ ID NO:), L1 (SEQ ID NO:), A8 (SEQ ID NO:), and A12 (SEQ ID NO:), respectively. The CDR regions (CDR1, CDR2 and CDR3) of each light chain have also been identified as noted in regard to Figures 1A-1G above.

Figures 3A-3G depict the κ light chain nucleic acid sequences of the Fab molecule clones 1:5 (SEQ ID NO:), 1:7 (SEQ ID NO:), 1:11 (SEQ ID NO:), L3 (SEQ ID NO:), L1 (SEQ ID NO:), A8 (SEQ ID NO:), and A12 (SEQ ID NO:), respectively.

Figures 4A-4G depict the γ 1 heavy chain nucleic acid sequences of the Fab molecule clones 1:5 (SEQ ID NO:), 1:7 (SEQ ID NO:), 1:11 (SEQ ID NO:), L3 (SEQ ID NO:), L1 (SEQ ID NO:), A8 (SEQ ID NO:), and A12 (SEQ ID NO:), respectively.

Figure 5 depicts the results of a neutralization assay conducted as described in Example 12 wherein the ability of Fab molecules expressed from the Fab molecule clones 1:5, 1:7, 1:11 and L3 to block the binding of HCV E2 polypeptide to target cells was assessed.

Detailed Description of the Invention

The practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al. *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *Nucleic Acid Hybridization* (B. Hames & S. Higgins, eds., 1985); *Transcription and Translation* (B. Hames & S. Higgins, eds., 1984); *Animal Cell Culture* (R. Freshney, ed., 1986); Perbal, *A Practical*

Guide to Molecular Cloning (1984); *Fundamental Virology*, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.).

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise.

A. Definitions

10 In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

As used herein, the terms "hepatitis C virus," or "HCV" describe the virus in a generic
15 manner, and as such the terms are not limiting to any particular HCV viral sequence or isolate. In this regard, there are 6 distinct genotypes of HCV with 11 distinct subtypes which have been identified based on phylogenetic analyses (Houghton, M. (1996) "Hepatitis
20 C Viruses," *Fields Virology*, 3d Edition, Fields et al. eds., Lippincott-Raven Publishers, Philadelphia PA.; Simmonds et al., *J. Gen. Virol.* (1993) 74:2391-2399). Further, comparison of genomic nucleotide sequences from different HCV isolates around the world establish
25 that HCV is highly heterogenous, with a range of sequence diversity among 74 isolates. Thus, the terms "hepatitis C virus," and "HCV" as used herein will generically encompass all such isolates.

The terms "an antigen derived from an E1
30 polypeptide," an "E1 polypeptide antigen" and "an HCV E1 antigen" are used interchangeably herein and encompass molecules from an HCV E1 region. The term "polypeptide," as used herein, refers to a polymer of amino acids and does not refer to a specific length of
35 the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or

exclude post expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. E1 polypeptides antigens can be physically derived from the HCV E1 region or produced recombinantly or synthetically, based on the known sequence. The mature E1 region of HCV1 begins at approximately amino acid 192 of the polyprotein and continues to approximately amino acid 383.

10 A polypeptide or amino acid sequence "derived from" a designated HCV region refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 3-5 amino acids, preferably at least 4-7 amino acids, more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence. This terminology also includes a polypeptide expressed from a designated HCV region.

 The terms "an antigen derived from an E2 polypeptide," an "E2 polypeptide antigen" and "an HCV E2 antigen" are used interchangeably herein and encompass molecules from an HCV E2 region. Such molecules can be physically derived from the region or produced recombinantly or synthetically, based on the known sequence. The mature E2 region of HCV1 is believed to begin at approximately amino acid 384-385.

30 For purposes of the present invention, HCV E1 and E2 polypeptides are defined with respect to the amino acid number of the polyprotein encoded by the genome of HCV1, with the initiator methionine being designated position 1. However, it should be noted that an antigen from an "E1 polypeptide" or an "E2 polypeptide" is not limited to polypeptides having an exact HCV1 sequence. Indeed, the HCV genome is in a

state of constant flux and contains several variable domains which exhibit relatively high degrees of variability between isolates. It is readily apparent that the terms encompass antigens from E1 and E2 polypeptides from any of the various HCV isolates including isolates having any of the 6 genotypes of HCV described in Simmonds et al., *J Gen Virol* (1993) 74:2391-2399). In this regard, the corresponding E1 or E2 regions in a heterologous HCV isolate can be readily determined by aligning sequences from the two isolates in a manner that brings the sequences into maximum alignment. This can be performed with any of a number of computer software packages, such as ALIGN 1.0, available from the University of Virginia, Department of Biochemistry (Attn: Dr. William R. Pearson). See, Pearson et al., *Proc Natl Acad Sci USA* (1988) 85:2444-2448.

Additionally, the terms "E1 polypeptide antigen" and "E2 polypeptide antigen" encompass polypeptides which include modifications to the native sequence, such as internal deletions, additions and substitutions (generally conservative in nature). These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through naturally occurring mutational events.

An "E1/E2 complex" refers to a complex of the E1 and E2 polypeptides described above. The mode of association of E1 and E2 in such a complex is immaterial. Indeed, such a complex may form spontaneously simply by mixing E1 and E2 polypeptides which have been produced individually. Similarly, when co-expressed and secreted, E1 and E2 polypeptides can form a complex spontaneously in the media. Formation of an "E1/E2 complex" is readily determined using standard protein detection techniques such as polyacrylamide gel electrophoresis and immunological techniques such as immunoprecipitation.

The term "antibody" encompasses monoclonal antibody preparations, as well as preparations including hybrid antibodies, altered antibodies, F(ab'), fragments, F(ab) molecules, Fv fragments, single domain antibodies, chimeric antibodies and functional fragments thereof which exhibit immunological binding properties of the parent antibody molecule.

As used herein, the term "monoclonal antibody" refers to an antibody composition having a homogeneous antibody population. The term is not limited by the manner in which it is made. The term encompasses whole immunoglobulin molecules, as well as Fab molecules, F(ab'), fragments, Fv fragments, and other molecules that exhibit immunological binding properties of the parent monoclonal antibody molecule. The term "recombinant monoclonal antibody" is defined herein as a monoclonal antibody that has been produced by expression of a recombinant polynucleotide.

The term "antigen-binding site," or "binding portion" refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable ("V") regions of the heavy ("H") and light ("L") chains. Three highly divergent stretches within the V regions of the heavy and light chains, referred to as "hypervariable regions," are interposed between more conserved flanking stretches known as "framework regions," or "FRs". Thus the term "FR" refers to amino acid sequences which are naturally found between, and adjacent to, hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary

to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as "complementarity-determining regions," or "CDRs."

5 As used herein, the terms "immunological binding," and "immunological binding properties" refer to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The
10 strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected
15 polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the
20 affinity of the interaction, and geometric parameters that equally influence the rate in both directions. Thus, both the "on rate constant" (K_{on}) and the "off rate constant" (K_{off}) can be determined by calculation of the concentrations and the actual rates of
25 association and dissociation. The ratio of K_{off}/K_{on} enables cancellation of all parameters not related to affinity, and is thus equal to the dissociation constant K_d . See, generally, Davies et al. (1990) *Annual Rev Biochem* 59:439-473.

30 A number of therapeutically useful molecules are known in the art which comprise antigen-binding sites that are capable of exhibiting immunological binding properties of an antibody molecule. One such molecule is a Fab molecule which comprises a
35 heterodimer that includes an intact antigen-binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the

"F(ab')₂" fragment which comprises both antigen-binding sites. An "Fv" fragment can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent V_H::V_L heterodimer including an antigen-binding site which retains much of the antigen recognition and binding capabilities of the native antibody molecule. Inbar et al. (1972) *Proc. Nat. Acad. Sci. USA* 69:2659-2662; Hochman et al. (1976) *Biochem* 15:2706-2710; and Ehrlich et al. (1980) *Biochem* 19:4091-4096.

A polypeptide molecule, or amino acid sequence "derived from" a designated Fab molecule or Fab nucleic acid sequence refers to a polypeptide having an amino acid sequence identical to that of a Fab polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 3-5 amino acids, preferably at least 4-7 amino acids, more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids.

A single chain Fv ("sFv") polypeptide molecule is a covalently linked V_H::V_L heterodimer which is expressed from a gene fusion including V_H- and V_L-encoding genes linked by a peptide-encoding linker. Huston et al. (1988) *Proc Nat Acad Sci USA* 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated, but chemically separated, light and heavy polypeptide chains from an antibody V region into an Sfv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Patent Nos. 5,091,513 and 5,132,405, to Huston et al.; and 4,946,778, to Ladner et al.

Each of the above-described molecules includes a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain FR set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term "CDR set" refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are denoted as "CDR1," "CDR2," and "CDR3," respectively. An antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region.

As used herein, the term "FR set" refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRs.

By "purified" and "isolated" is meant, when referring to a polypeptide or nucleotide sequence, that the indicated molecule is present in the substantial absence of other biological macromolecules of the same type. The terms "purified" and "isolated" as used herein preferably mean at least 75% by weight, more preferably at least 85% by weight, more preferably still at least 95% by weight, and most preferably at least 98% by weight, of biological macromolecules of the same type are present. An "isolated nucleic acid molecule which encodes a particular polypeptide" refers to a nucleic acid molecule which is substantially free of other nucleic acid molecules that do not encode the subject polypeptide; however, the molecule may include some additional bases or moieties which do not deleteriously affect the basic characteristics of the

composition. Thus, for example, an isolated nucleic acid molecule which encodes the binding portion of a particular heavy chain variable region of an antibody consists essentially of the nucleotide coding sequence
5 for the subject binding portion (e.g., the CDR set interposed between the FR set).

"Homology" refers to the percent of identity between two polynucleotide or polypeptide moieties. The correspondence between two or more sequences can
10 be determined by techniques known in the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide molecules. Alternatively, homology can be determined by hybridization of polynucleotides under
15 conditions which form stable duplexes between homologous regions (for example, those which would be used prior to S_1 digestion), followed by digestion with single-stranded specific nuclease(s), followed by size determination of the digested fragments. Two DNA
20 or polypeptide sequences are "substantially homologous" when at least about 60% (preferably at least about 80%, and most preferably at least about 90%) of the nucleotides or amino acids match over a defined length of the molecule.

25 The terms "recombinant DNA molecule," or "recombinant nucleic acid molecule" are used herein to refer to a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with
30 all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature. Thus, the term encompasses "synthetically derived" nucleic acid
35 molecules.

The term "nucleic acid molecule" as used herein refers to a polymeric form of nucleotides of

any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule and thus includes double- and single-stranded DNA and RNA. It also includes known types of modifications, for example, labels which are known in the art, methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, for example proteins (including for e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide.

A "coding sequence" is a nucleic acid molecule which is translated into a polypeptide, usually via mRNA, when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence may be determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A coding sequence can include, but is not limited to, CDNA, and recombinant nucleotide sequences.

"Control sequence" refers to nucleic acid sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal

binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include, at a minimum, all components whose presence is necessary for expression of a coding sequence, and may also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

"Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

As used herein, the term "expression cassette" refers to a molecule comprising at least one coding sequence operably linked to a control sequence which includes all nucleotide sequences required for the transcription of cloned copies of the coding sequence and the translation of the mRNAs in an appropriate host cell. Such expression cassettes can be used to express eukaryotic genes in a variety of hosts such as bacteria, blue-green algae, plant cells, yeast cells, insect cells and animal cells. Under the invention, expression cassettes can include, but are not limited to, cloning vectors, specifically designed plasmids, viruses or virus particles. The cassettes may further include an origin of replication for autonomous replication in host cells, selectable markers, various restriction sites, a potential for high copy number and strong promoters.

By "vector" is meant any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus etc., which is capable of replication when associated with the proper control

elements and which can transfer gene sequences between cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

"Recombinant host cells", "host cells,"
5 "cells," "cell cultures," and other such terms denote, for example, microorganisms, insect cells, and mammalian cells, that can be, or have been, used as recipients for recombinant vector or other transfer DNA, and include the progeny of the original cell
10 which has been transformed. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation. Examples
15 for mammalian host cells include Chinese hamster ovary (CHO) and monkey kidney (COS) cells.

Specifically, as used herein, "cell line," refers to a population of cells capable of continuous or prolonged growth and division in vitro. Often,
20 cell lines are clonal populations derived from a single progenitor cell. It is further known in the art that spontaneous or induced changes can occur in karyotype during storage or transfer of such clonal populations. Therefore, cells derived from the cell
25 line referred to may not be precisely identical to the ancestral cells or cultures, and the cell line referred to includes such variants. The term "cell lines" also includes immortalized cells.

"Transformation", as used herein, refers to
30 the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion, for example, direct uptake, transduction, f-mating or electroporation. The exogenous polynucleotide may be maintained as a non-integrated
35 vector, for example, a plasmid, or alternatively, may be integrated into the host genome.

B. General Methods

The present invention is based on the generation of novel cross-genotype reactive human monoclonal antibody molecules specific to the HCV E2 envelope glycoprotein. The monoclonal antibodies are obtained using a combinatorial antibody library constructed from a nonimmunized source, and are useful in the prevention, therapy and diagnosis of HCV infection in mammalian subjects. More particularly, the monoclonal antibodies are obtained from combinatorial libraries expressing Fab molecules on the surface of filamentous DNA bacteriophage using antigen selection techniques.

Preparation of Combinatorial Libraries

Combinatorial libraries for the purposes of the present invention can be constructed using known techniques, such as those described by Chanock et al. (1993) *Infect Agents Dis* 2:118-131 and Barbas, III et al. (1995) *Methods: Comp. Meth Enzymol* 8:94-103. Antibody-producing cells can be obtained from an unimmunized, HCV-infected donor from, e.g., plasma, serum, spinal fluid, lymph fluid, the external sections of the respiratory, intestinal and genitourinary tracts, tears, saliva, milk, white blood cells and myelomas. Preferably, the antibody-producing cell source is lymphocytes that have been obtained from a bone marrow or peripheral blood sample of an unimmunized subject.

Lymphocytes can be obtained from the sample and total RNA isolated and extracted using known methods. See, e.g., Chomczynski et al. (1987) *Anal Biochem* 162:156-159. The RNA can be reverse-transcribed into first strand cDNA using oligo-dT priming. The DNA encoding immunoglobulin heavy (Fd) and light chain fragments can be amplified using the polymerase chain reaction (PCR) to provide all of the

genetic material necessary to produce Fab antigen-binding molecules. Saiki, et al. (1986) *Nature* 324:163, Scharf et al. (1986) *Science* 233:1076-1078 and U.S. Patent Nos. 4,683,195 and 4,683,202. In

5 conducting the PCR amplification, a number of known primers can preferably be used to select for γ 1 heavy chain and κ light chain sequences. Persson et al. (1991) *Proc Natl Acad Sci USA* 88:2432-2436, Kang et al. (1991) *Methods: Comp. Meth Enzymol* 2:111-118. The

10 PCR products are pooled separately into heavy and light chain DNA preparations, and then purified, for example, using gel electrophoresis. The purified heavy and light chain DNA molecules are then digested with suitable restriction enzymes, and the digested

15 products purified and ligated into a suitable phagemid vector system. Yang et al. (1995) *J Mol Biol* 254:392-403, Barbas, III et al. (1995) *Methods: Comp. Meth Enzymol* 8:94-103, Barbas, III et al. (1991) *Proc Natl Acad Sci USA* 88:7978-7982. A number of suitable

20 phagemid vector systems are known in the art; however, a particularly preferred vector for use herein is the pComb3H vector which has been previously described. Barbas, III et al. (1995), *supra*. When the Pcomb3h phagemid vector is used, heavy chain DNA is cloned

25 into the subject phagemid adjacent to, and upstream of, the sequence for the C-terminal anchorage domain of the phagemid coat protein III (cpIII). The cpIII protein is an integral membrane protein, and thus serves as a membrane anchor for the Fab assembly.

30 The vectors generally include selectable markers known in the art. For example, the Pcomb3h phagemid vector contains the bacterial ampicillin resistance gene (B-lactamase). The vector will also include appropriate first and second leader sequences,

35 respectively arranged upstream of the insertion sites for the heavy and light chain coding sequences, whereby expression products from the heavy and light

chain coding regions are targeted to the periplasm when produced in a suitable host cell. In Pcomb3h, these leader sequences are *pelB* sequences, *omp A* sequences or combinations thereof.

5 The phagemid vector system containing the human immunoglobulin DNA is then introduced into a suitable bacterial host cell (for example using electrophoresis), wherein the phagemid expresses a heavy chain-*cpiii* fusion polypeptide and a light chain
10 polypeptide, each of which are targeted to the periplasm of the host cell by their associated leader sequences. The transfected bacterial host cell containing the phagemid vector is selected by growth in a suitable medium containing a selective agent
15 corresponding to the selectable marker of the phagemid vector (e.g., ampicillin).

 Rescue of the phagemid DNA is conducted using known techniques. In particular, the transfected host cell is infected with a helper phage
20 which encodes a number of expression products necessary *in trans* for packaging the phagemid DNA into recombinant virus particles. Single-stranded copies of the phagemid DNA are thus packaged into viral particles which, upon leaving the host cell,
25 incorporate phage *cpVIII* molecules and are capped by a limited number of phage *cpiii* molecules--some of which *cpviii* and *cpiii* molecules are linked to Fab molecules. Recombinant phage particles displaying Fab molecules (termed "phage-Fabs") contain the
30 corresponding heavy and light chain genes within the packaged genome.

 When the above technique is practiced using an initial library of phagemids, the rescue process generates a library of recombinant phage which display
35 Fab molecules (a phage display library). The rescue process further results in amplification of the initial library, such that multiple copies of each

recombinant phage clone (along with each set of immunoglobulin heavy and light chain binding portions) are generated. The phage display library is then "panned" against HCV E2 antigen to select for Fab molecules which are capable of selectively binding to that antigen. More particularly, the panning procedure can be conducted by applying a suspension containing the phage display library onto HCV E2 antigen that has been immobilized to a plastic reaction vessel according to known methods. Burton et al. (1991) *Proc Natl Acad Sci USA* 88:10134-10137. After incubation under suitable binding conditions, non-specifically bound phage particles are removed by repeated washings. The resulting HCV E2-antigen specific phage-Fabs are then eluted from the insoluble antigen using low Ph, or in the presence of excess soluble E2 antigen. The panning procedure is repeated several times, wherein bacterial host cells are infected by the eluted phage after each round of panning to propagate phage-Fab clones for each subsequent round of panning. Samuelsson et al. (1995) *Virology* 207:495-502.

In the present invention, the panning procedure was specifically developed to select for highly potent, cross-genotype reactive Fab molecules specific for HCV E2 antigen. In particular, the genotype of serum HCV of the unimmunized, HCV-infected human subject from which the antibody-producing cells were obtained was determined using known methods. Widell et al. (1994) *J Med Virol* 44:272-279. Selection for strain cross-reactivity was provided by experimental design, wherein the panning procedure was conducted using HCV E2 antigen derived from a different HCV genotype than that of the HCV from the infected human donor. Furthermore, the E2 antigen used in the panning procedures was selected so as to

provide HCV E2 antigen in substantially the same conformation as expected for that antigen *in vivo*.

Two different recombinant HCV envelope protein preparations were used to provide the
5 selecting antigen in the above-described panning procedure, a "conformational" CHO E2 molecule, and a CHO E1/E2 complex. The conformational E2 molecule was constructed, expressed and secreted from recombinant CHO cells as previously described in Spaete et al.
10 (1992) *Virology* 188:819-830, then purified using known methods (Rosa et al. (1996) *Proc Natl Acad Sci USA* 93:1759-1763). A recombinant complexed E1/E2 preparation was constructed and expressed from recombinant CHO cells as described in Spaete et al.
15 (1992) *supra*, then purified using known methods (Choo et al. (1994) *Proc Natl Acad Sci USA* 91:1294-1298). Once purified, the selecting antigens were immobilized to a plastic reaction vessel as described above.

Individual clones exhibiting superior
20 binding affinity for the selecting antigen were selected, and expressed by growing infected host cells in the selective medium until a suitable volume of cells was reached. The bacterial host cells were pelleted and then resuspended in medium. After
25 suitable incubation, the cells were spun down, and the periplasmic content released by freeze-thawing techniques. After the bacterial debris was removed by centrifugation, the Fab-containing supernatant was transferred to suitable containers, and stored for
30 future use.

Once the selected Fabs are expressed, binding characteristics of the selected Fab molecules can be determined. In particular, the affinity of the Fab molecules for HCV E2 antigen was determined herein
35 using an inhibition ELISA technique. See, e.g., Persson et al. (1991) *Proc Natl Acad Sci USA* 88:2432-2436, Rath et al. (1988) *J Immun Methods* 106:245-249.

Clones that expressed Fab molecules of high potency (e.g., an affinity of at least about $1 \times 10^7 \text{ M}^{-1}$, and preferably at least about $1.7 \times 10^7 \text{ M}^{-1}$ as determined by inhibition ELISA) were identified for sequencing.

5 Phage (plasmid) DNA from clones which exhibited high potency binding in the panning selection process was isolated, and single stranded DNA was obtained by PCR using primers (one of which, e.g., is biotinylated at the 5' end) that hybridize
10 upstream and downstream of the immunoglobulin cloning regions. After PCR, single stranded DNA was obtained by denaturing the DNA under alkaline conditions, and absorbing biotinylated DNA strands onto a solid support. Dideoxy sequencing reactions were performed
15 according to known methods (Sanger et al. (1977) *Proc Natl Acad Sci USA* 74:5463-5467) using labeled primers hybridizing 3' of the junction between the variable and constant regions. Kabat et al., in *Sequences of Proteins of Immunological Interest*, 4th ed., (U.S. Dept. of Health and Human Services, U.S. Government
20 Printing Office, 1987). The reaction products were run on an automated sequencer (for example, A.L.F. available from Pharmacia Biotech). The nucleic acid sequence information thus obtained was analyzed to
25 provide coding sequences for the heavy chain and light chain portions of the selected monoclonal Fab molecules. Multiple copies of the same clones were identified by comparisons of sequence data. Further, the deduced amino acid sequences were obtained using
30 known methods.

 Using the above nucleic acid sequence information, coding sequences for the Fab molecules can also be produced synthetically using known methods. Nucleotide sequences can be designed with
35 the appropriate codons for the particular amino acid sequence desired. In general, one will select preferred codons for the intended host in which the

sequences will be expressed. The complete sequences are generally assembled from overlapping oligonucleotides prepared by standard methods and assembled into complete coding sequences. See, e.g.,
5 Edge (1981) *Nature* 292:756; Nambair et al. (1984) *Science* 223:1299; Jay et al. (1984) *J. Biol. Chem.* 259:6311.

Expression Systems

10 Once the coding sequences for the heavy and light chain portions of the Fab molecules are isolated or synthesized, they can be cloned into any suitable vector or replicon for expression, for example, bacterial, mammalian, yeast and viral expression
15 systems can be used. Numerous cloning vectors are known to those of skill in the art and are described below. The selection of an appropriate cloning vector is a matter of choice.

20 i. Expression in Bacterial Cells

Bacterial expression systems can be used to produce the Fab molecules. Control elements for use in bacterial systems include promoters, optionally containing operator sequences, and ribosome binding
25 sites. Useful promoters include sequences derived from sugar metabolizing enzymes, such as galactose, lactose (*lac*) and maltose. Additional examples include promoter sequences derived from biosynthetic enzymes such as tryptophan (*trp*), the β -lactamase
30 (*bla*) promoter system, bacteriophage λ PL, and T7. In addition, synthetic promoters can be used, such as the *tac* promoter. The β -lactamase and lactose promoter systems are described in Chang et al., *Nature* (1978) 275:615, and Goeddel et al., *Nature* (1979) 281: 544;
35 the alkaline phosphatase, tryptophan (*trp*) promoter system are described in Goeddel et al., *Nucleic Acids Res.* (1980) 8:4057 and EP 36,776 and hybrid promoters

such as the *tac* promoter is described in U.S. Patent No. 4,551,433 and deBoer et al., *Proc. Natl. Acad. Sci. USA* (1983) 80:21-25. However, other known bacterial promoters useful for expression of eukaryotic proteins are also suitable. A person skilled in the art would be able to operably ligate such promoters to the Fab molecules for example, as described in Siebenlist et al., *Cell* (1980) 20:269, using linkers or adapters to supply any required restriction sites. Promoters for use in bacterial systems also generally contain a Shine-Dalgarno (SD) sequence operably linked to the DNA encoding the Fab molecule. For prokaryotic host cells that do not recognize and process the native polypeptide signal sequence, the signal sequence can be substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, Ipp, or heat stable enterotoxin II leaders. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria.

The foregoing systems are particularly compatible with *Escherichia coli*. However, numerous other systems for use in bacterial hosts including Gram-negative or Gram-positive organisms such as *Bacillus spp.*, *Streptococcus spp.*, *Streptomyces spp.*, *Pseudomonas* species such as *P. aeruginosa*, *Salmonella typhimurium*, or *Serratia marcescans*, among others. Methods for introducing exogenous DNA into these hosts typically include the use of CaCl_2 or other agents, such as divalent cations and DMSO. DNA can also be introduced into bacterial cells by electroporation, nuclear injection, or protoplast fusion as described generally in Sambrook et al. (1989), cited above. These examples are illustrative rather than limiting. Preferably, the host cell should secrete minimal amounts of proteolytic enzymes. Alternatively, in

vitro methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

Prokaryotic cells used to produce the Fab molecules of this invention are cultured in suitable media, as described generally in Sambrook et al.,
5 cited above.

ii. Expression in Yeast Cells

Yeast expression systems can also be used to
10 produce the subject Fab molecules. Expression and transformation vectors, either extrachromosomal replicons or integrating vectors, have been developed for transformation into many yeasts. For example, expression vectors have been developed for, among
15 others, the following yeasts: *Saccharomyces cerevisiae*, as described in Hinnen et al., *Proc. Natl. Acad. Sci. USA* (1978) 75:1929; Ito et al., *J. Bacteriol.* (1983) 153:163; *Candida albicans* as described in Kurtz et al., *Mol. Cell. Biol.* (1986)
20 6:142; *Candida maltosa*, as described in Kunze et al., *J. Basic Microbiol.* (1985) 25:141; *Hansenula polymorpha*, as described in Gleeson et al., *J. Gen. Microbiol.* (1986) 132:3459 and Roggenkamp et al., *Mol. Gen. Genet.* (1986) 202:302; *Kluyveromyces fragilis*, as
25 described in Das et al., *J. Bacteriol.* (1984) 158:1165; *Kluyveromyces lactis*, as described in De Louvencourt et al., *J. Bacteriol.* (1983) 154:737 and Van den Berg et al., *Bio/Technology* (1990) 8:135; *Pichia guillerimondii*, as described in Kunze et al.,
30 *J. Basic Microbiol.* (1985) 25:141; *Pichia pastoris*, as described in Cregg et al., *Mol. Cell. Biol.* (1985) 5:3376 and U.S. Patent Nos. 4,837,148 and 4,929,555; *Schizosaccharomyces pombe*, as described in Beach and Nurse, *Nature* (1981) 300:706; and *Yarrowia lipolytica*,
35 as described in Davidow et al., *Curr. Genet.* (1985) 10:380 and Gaillardin et al., *Curr. Genet.* (1985) 10:49, *Aspergillus* hosts such as *A. nidulans*, as

described in Ballance et al., *Biochem. Biophys. Res. Commun.* (1983) 112:284-289; Tilburn et al., *Gene* (1983) 26:205-221 and Yelton et al., *Proc. Natl. Acad. Sci. USA* (1984) 81:1470-1474, and *A. niger*, as
5 described in Kelly and Hynes, *EMBO J.* (1985) 4:475479; *Trichoderma reesia*, as described in EP 244,234, and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, as described in WO 91/00357.

10 Control sequences for yeast vectors are known and include promoter regions from genes such as alcohol dehydrogenase (ADH), as described in EP 284,044, enolase, glucokinase, glucose-6-phosphate isomerase, glyceraldehyde-3-phosphate-dehydrogenase
15 (GAP or GAPDH), hexokinase, phosphofructokinase, 3-phosphoglycerate mutase, and pyruvate kinase (PyK), as described in EP 329,203. The yeast *PHO5* gene, encoding acid phosphatase, also provides useful promoter sequences, as described in Myanohara et al.,
20 *Proc. Natl. Acad. Sci. USA* (1983) 80:1. Other suitable promoter sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase, as described in Hitzeman et al., *J. Biol. Chem.* (1980) 255:2073, or other glycolytic enzymes, such as
25 pyruvate decarboxylase, triosephosphate isomerase, and phosphoglucose isomerase, as described in Hess et al., *J. Adv. Enzyme Reg.* (1968) 7:149 and Holland et al., *Biochemistry* (1978) 17:4900. Inducible yeast promoters having the additional advantage of
30 transcription controlled by growth conditions, include from the list above and others the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein,
35 glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast

expression are further described in Hitzeman, EP 073,657. Yeast enhancers also are advantageously used with yeast promoters. In addition, synthetic promoters which do not occur in nature also function as yeast promoters. For example, upstream activating sequences (UAS) of one yeast promoter may be joined with the transcription activation region of another yeast promoter, creating a synthetic hybrid promoter. Examples of such hybrid promoters include the ADH regulatory sequence linked to the GAP transcription activation region, as described in U.S. Patent Nos. 4,876,197 and 4,880,734. Other examples of hybrid promoters include promoters which consist of the regulatory sequences of either the *ADH2*, *GAL4*, *GAL10*, or *PHO5* genes, combined with the transcriptional activation region of a glycolytic enzyme gene such as *GAP* or *PyK*, as described in EP 164,556. Furthermore, a yeast promoter can include naturally occurring promoters of non-yeast origin that have the ability to bind yeast RNA polymerase and initiate transcription.

Other control elements which may be included in the yeast expression vectors are terminators, for example, from *GAPDH* and from the enolase gene, as described in Holland et al., *J. Biol. Chem.* (1981) 256:1385, and leader sequences which encode signal sequences for secretion. DNA encoding suitable signal sequences can be derived from genes for secreted yeast proteins, such as the yeast invertase gene as described in EP 012,873 and JP 62,096,086 and the α -factor gene, as described in U.S. Patent Nos. 4,588,684, 4,546,083 and 4,870,008; EP 324,274; and WO 89/02463. Alternatively, leaders of non-yeast origin, such as an interferon leader, also provide for secretion in yeast, as described in EP 060,057.

Methods of introducing exogenous DNA into yeast hosts are well known in the art, and typically include either the transformation of spheroplasts or of

intact yeast cells treated with alkali cations. Transformations into yeast can be carried out according to the method described in Van Solingen et al., *J. Bact.* (1977) 130:946 and Hsiao et al., *Proc. Natl. Acad. Sci. USA* (1979) 76:3829. However, other methods for introducing DNA into cells such as by nuclear injection, electroporation, or protoplast fusion may also be used as described generally in Sambrook et al., cited above.

For yeast secretion the native polypeptide signal sequence may be substituted by the yeast invertase, α -factor, or acid phosphatase leaders. The origin of replication from the 2μ plasmid origin is suitable for yeast. A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid described in Kingsman et al., *Gene* (1979) 7:141 or Tschemper et al., *Gene* (1980) 10:157. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* Gene.

For intracellular production of the present polypeptides in yeast, a sequence encoding a yeast protein can be linked to a coding sequence for the Fab molecule to produce a fusion protein that can be cleaved intracellularly by the yeast cells upon expression. An example, of such a yeast leader sequence is the yeast ubiquitin gene.

30 iii. Expression in Insect Cells

The Fab molecules can also be produced in insect expression systems. For example, baculovirus expression vectors (BEVs) are recombinant insect viruses in which the coding sequence for a foreign gene to be expressed is inserted behind a baculovirus promoter in place of a viral gene, e.g., polyhedrin,

as described in Smith and Summers, U.S. Pat. No.,
4,745,051.

5 An expression construct herein includes a
DNA vector useful as an intermediate for the infection
or transformation of an insect cell system, the vector
generally containing DNA coding for a baculovirus
transcriptional promoter, optionally but preferably,
followed downstream by an insect signal DNA sequence
capable of directing secretion of a desired protein,
10 and a site for insertion of the foreign gene encoding
the foreign protein, the signal DNA sequence and the
foreign gene being placed under the transcriptional
control of a baculovirus promoter, the foreign gene
herein being the coding sequence of the Fab molecule.

15 The promoter for use herein can be a
baculovirus transcriptional promoter region derived
from any of the over 500 baculoviruses generally
infecting insects, such as, for example, the Orders
Lepidoptera, Diptera, Orthoptera, Coleoptera and
20 Hymenoptera including, for example, but not limited to
the viral DNAs of *Autographa californica* MNPV, *Bombyx
mori* NPV, *Trichoplusia ni* MNPV, *Rachlplusia ou* MNPV or
Galleria mellonella MNPV, *Aedes aegypti*, *Drosophila
melanogaster*, *Spodoptera frugiperda*, and *Trichoplusia
25 ni*. Thus, the baculovirus transcriptional promoter
can be, for example, a baculovirus immediate-early
gene IEI or IEN promoter; an immediate-early gene in
combination with a baculovirus delayed-early gene
promoter region selected from the group consisting of
30 a 39K and a *HindIII* fragment containing a
delayed-early gene; or a baculovirus late gene
promoter. The immediate-early or delayed-early
promoters can be enhanced with transcriptional
enhancer elements.

35 Particularly suitable for use herein is the
strong polyhedrin promoter of the baculovirus, which
directs a high level of expression of a DNA insert, as

described in Friesen et al. (1986) "The Regulation of
Baculovirus Gene Expression" in: THE MOLECULAR BIOLOGY
OF BACULOVIRUSES (W.Doerfler, ed.); EP 127,839 and EP
155,476; and the promoter from the gene encoding the
5 p10 protein, as described in Vlak et al., J. Gen.
Viro1. (1988) 69:765-776.

The plasmid for use herein usually also
contains the polyhedrin polyadenylation signal, as
described in Miller et al., Ann. Rev. Microbiol.
10 (1988) 42:177 and a procaryotic ampicillin-resistance
(amp) gene and an origin of replication for selection
and propagation in *E. coli*. DNA encoding suitable
signal sequences can also be included and is generally
derived from genes for secreted insect or baculovirus
15 proteins, such as the baculovirus polyhedrin gene, as
described in Carbonell et al., Gene (1988) 73:409, as
well as mammalian signal sequences such as those
derived from genes encoding human α -interferon as
described in Maeda et al., Nature (1985) 315:592-594;
20 human gastrin-releasing peptide, as described in
Lebacqz-Verheyden et al., Mol. Cell. Biol. (1988)
8:3129; human IL-2, as described in Smith et al.,
Proc. Natl. Acad. Sci. USA (1985) 82:8404; mouse IL-3,
as described in Miyajima et al., Gene (1987) 58:273;
25 and human glucocerebrosidase, as described in Martin
et al., DNA (1988) 7:99.

Numerous baculoviral strains and variants
and corresponding permissive insect host cells from
hosts such as *Spodoptera frugiperda* (caterpillar),
30 *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito),
Drosophila melanogaster (fruitfly), and *Bombyx mori*
host cells have been identified and can be used
herein. See, for example, the description in Luckow
et al., Bio/Technology (1988) 6:47-55, Miller et al.,
35 in GENETIC ENGINEERING (Setlow, J.K. et al. eds.),
Vol. 8 (Plenum Publishing, 1986), pp. 277-279, and
Maeda et al., Nature (1985) 315:592-594. A variety of

such viral strains are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV. Such viruses may be used as the virus for transfection of host cells such as

5 *Spodoptera frugiperda* cells.

Other baculovirus genes in addition to the polyhedrin promoter may be employed in a baculovirus expression system. These include immediate-early (alpha), delayed-early (beta), late (gamma), or very

10 late (delta), according to the phase of the viral infection during which they are expressed. The expression of these genes occurs sequentially, probably as the result of a "cascade" mechanism of transcriptional regulation. Thus, the immediate-early

15 genes are expressed immediately after infection, in the absence of other viral functions, and one or more of the resulting gene products induces transcription of the delayed-early genes. Some delayed-early gene products, in turn, induce transcription of late genes,

20 and finally, the very late genes are expressed under the control of previously expressed gene products from one or more of the earlier classes. One relatively well defined component of this regulatory cascade is IEI, a preferred immediate-early gene of *Autographa californica* nuclear polyhedrosis virus (AcMNPV). IEI

25 is pressed in the absence of other viral functions and encodes a product that stimulates the transcription of several genes of the delayed-early class, including the preferred 39K gene, as described in Guarino and

30 Summers, *J. Virol.* (1986) 57:563-571 and *J. Virol.* (1987) 61:2091-2099 as well as late genes, as described in Guanno and Summers, *Virol.* (1988) 162:444-451.

Immediate-early genes as described above can

35 be used in combination with a baculovirus gene promoter region of the delayed-early category. Unlike the immediate-early genes, such delayed-early genes

require the presence of other viral genes or gene products such as those of the immediate-early genes. The combination of immediate-early genes can be made with any of several delayed-early gene promoter regions such as 39K or one of the delayed-early gene promoters found on the *HindIII* fragment of the baculovirus genome. In the present instance, the 39 K promoter region can be linked to the foreign gene to be expressed such that expression can be further controlled by the presence of IEI, as described in L. A. Guarino and Summers (1986a), cited above; Guarino & Summers (1986b) *J. Virol.* (1986) 60:215-223, and Guarino et al. (1986c) *J. Virol.* (1986) 60:224-229.

Additionally, when a combination of immediate-early genes with a delayed-early gene promoter region is used, enhancement of the expression of heterologous genes can be realized by the presence of an enhancer sequence in direct cis linkage with the delayed-early gene promoter region. Such enhancer sequences are characterized by their enhancement of delayed-early gene expression in situations where the immediate-early gene or its product is limited. For example, the hr5 enhancer sequence can be linked directly, in cis, to the delayed-early gene promoter region, 39K, thereby enhancing the expression of the cloned heterologous DNA as described in Guarino and Summers (1986a), (1986b), and Guarino et al. (1986).

The polyhedrin gene is classified as a very late gene. Therefore, transcription from the polyhedrin promoter requires the previous expression of an unknown, but probably large number of other viral and cellular gene products. Because of this delayed expression of the polyhedrin promoter, state-of-the-art BEVs, such as the exemplary BEV system described by Smith and Summers in, for example, U.S. Pat. No., 4,745,051 will express foreign genes only as a result of gene expression from the rest of

the viral genome, and only after the viral infection is well underway. This represents a limitation to the use of existing BEVs. The ability of the host cell to process newly synthesized proteins decreases as the baculovirus infection progresses. Thus, gene expression from the polyhedrin promoter occurs at a time when the host cell's ability to process newly synthesized proteins is potentially diminished for certain proteins such as human tissue plasminogen activator. As a consequence, the expression of secretory glycoproteins in BEV systems is complicated due to incomplete secretion of the cloned gene product, thereby trapping the cloned gene product within the cell in an incompletely processed form.

While it has been recognized that an insect signal sequence can be used to express a foreign protein that can be cleaved to produce a mature protein, the present invention can also be practiced with a mammalian signal sequence.

An exemplary insect signal sequence suitable herein is the sequence encoding for a Lepidopteran adipokinetic hormone (AKH) peptide. The AKH family consists of short blocked neuropeptides that regulate energy substrate mobilization and metabolism in insects. In a preferred embodiment, a DNA sequence coding for a Lepidopteran *Manduca sexta* AKH signal peptide can be used. Other insect AKH signal peptides, such as those from the Orthoptera *Schistocerca gregaria* locus can also be employed to advantage. Another exemplary insect signal sequence is the sequence coding for *Drosophila* cuticle proteins such as CPI, CP2, CP3 or CP4.

Currently, the most commonly used transfer vector that can be used herein for introducing foreign genes into AcNPV is pAc373. Many other vectors, known to those of skill in the art, can also be used herein. Materials and methods for baculovirus/insect cell

expression systems are commercially available in a kit form from companies such as Invitrogen (San Diego CA) ("MaxBac" kit). The techniques utilized herein are generally known to those skilled in the art and are
5 fully described in Summers and Smith, A MANUAL OF METHODS FOR BACULOVIRUS VECTORS AND INSECT CELL CULTURE PROCEDURES, Texas Agricultural Experiment Station Bulletin No. 1555, Texas A&M University (1987); Smith et al., *Mol. Cell. Biol.* (1983), and
10 Luckow and Summers (1989). These include, for example, the use of pVL985 which alters the polyhedrin start codon from ATG to ATT, and which introduces a *Bam*HI cloning site 32 basepairs downstream from the ATT, as described in Luckow and Summers, *Virology*
15 (1989) 17:31.

Thus, for example, for insect cell expression of the present polypeptides, the desired DNA sequence can be inserted into the transfer vector, using known techniques. An insect cell host can be
20 cotransformed with the transfer vector containing the inserted desired DNA together with the genomic DNA of wild type baculovirus, usually by cotransfection. The vector and viral genome are allowed to recombine resulting in a recombinant virus that can be easily
25 identified and purified. The packaged recombinant virus can be used to infect insect host cells to express a Fab molecule.

Other methods that are applicable herein are the standard methods of insect cell culture,
30 cotransfection and preparation of plasmids are set forth in Summers and Smith (1987), cited above. This reference also pertains to the standard methods of cloning genes into AcMNPV transfer vectors, plasmid DNA isolation, transferring genes into the AcMNPV
35 genome, viral DNA purification, radiolabeling recombinant proteins and preparation of insect cell culture media. The procedure for the cultivation of

viruses and cells are described in Volkman and Summers, *J. Virol.* (1975) 19:820-832 and Volkman et al., *J. Virol.* (1976) 19:820-832.

5 iv. Expression in Mammalian Cells

Mammalian expression systems can also be used to produce the Fab molecules. Typical promoters for mammalian cell expression include the SV40 early promoter, the CMV promoter, the mouse mammary tumor virus LTR promoter, the adenovirus major late promoter (Ad MLP), and the herpes simplex virus promoter, among others. Other non-viral promoters, such as a promoter derived from the murine metallothionein gene, will also find use in mammalian constructs. Mammalian expression may be either constitutive or regulated (inducible), depending on the promoter. Typically, transcription termination and polyadenylation sequences will also be present, located 3' to the translation stop codon. Preferably, a sequence for optimization of initiation of translation, located 5' to the Fab coding sequence, is also present. Examples of transcription terminator/polyadenylation signals include those derived from SV40, as described in Sambrook et al. (1989) *MOLECULAR CLONING: A LABORATORY*
10
15
20
25
30
35
MANUAL, 2d edition, (Cold Spring Harbor Press, Cold Spring Harbor, N.Y.). Introns, containing splice donor and acceptor sites, may also be designed into the constructs of the present invention.

Enhancer elements can also be used herein to increase expression levels of the mammalian constructs. Examples include the SV40 early gene enhancer, as described in Dijkema et al., *EMBO J.* (1985) 4: 761 and the enhancer/promoter derived from the long terminal repeat (LTR) of the Rous Sarcoma Virus, as described in Gorman et al., *Proc. Natl. Acad. Sci. USA* (1982b) 79:6777 and human cytomegalovirus, as described in Boshart et al., *Cell*

(1985) 41:521. A leader sequence can also be present which includes a sequence encoding a signal peptide, to provide for the secretion of the foreign protein in mammalian cells. Preferably, there are processing sites encoded between the leader fragment and the gene of interest such that the leader sequence can be cleaved either *in vivo* or *in vitro*. The adenovirus tripartite leader is an example of a leader sequence that provides for secretion of a foreign protein in mammalian cells.

There exist expression vectors that provide for the transient expression in mammalian cells of DNA encoding the Fab molecules. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of polypeptides encoded by cloned DNAs, as well as for the rapid screening of such polypeptides for desired biological or physiological properties. Once complete, the mammalian expression vectors can be used to transform any of several mammalian cells. Methods for introduction of heterologous polynucleotides into mammalian cells are known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. General aspects of mammalian cell host system transformations have been described by Axel in U.S. 4,399,216. A synthetic lipid particularly useful for polynucleotide transfection is N-[1-(2,3-dioleyloxy)propyl]-N,N,N-

trimethylammonium chloride, which is commercially available under the name Lipofectin® (available from BRL, Gaithersburg, MD), and is described by Felgner et al., *Proc. Natl. Acad. Sci. USA* (1987) 84:7413.

5 Mammalian cell lines available as hosts for expression are also known and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to, Chinese hamster ovary (CHO) cells, HeLa
10 cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), human embryonic kidney cells, baby hamster kidney cells, mouse sertoli cells, canine kidney cells, buffalo rat liver cells, human lung
15 cells, human liver cells, mouse mammary tumor cells, as well as others.

The mammalian host cells used to produce the Fab molecules of this invention may be cultured in a variety of media. Commercially available media such
20 as Ham's F10 (Sigma), Minimal Essential Medium ([MEM], Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ([DMEM], Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham and Wallace, *Meth. Enz.* (1979)
25 58:44, Barnes and Sato, *Anal. Biochem.* (1980) 102:255, U.S. Patent Nos. 4,767,704, 4,657,866, 4,927,762, or 4,560,655, WO 90/103430, WO 87/00195, and U.S. RE 30,985, may be used as culture media for the host cells. Any of these media may be supplemented as
30 necessary with hormones and/or other growth factors such as insulin, transferrin, or epidermal growth factor, salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine),
35 antibiotics (such as Gentamycin(tm) M drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar

range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, Ph, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

10 Preparing Specific Binding Molecules

Using the above techniques, a number of specific binding molecules that exhibit immunological binding affinity for HCV E2 antigen can be provided. In particular, depending on the expression system and host selected, soluble Fab specific binding molecules can be readily produced by growing host cells transformed by an expression vector described above under conditions whereby the heavy and light chain portions are expressed. Heterodimers comprising noncovalently associated heavy and light chains can be isolated from the host cells and purified. Since the present invention also provides for the optional secretion of the heavy and light chain polypeptides, the Fab heterodimers can be purified directly from the media. The selection of the appropriate growth conditions and recovery methods are within the skill of the art.

In addition, the Fab molecules of the present invention can be produced using conventional methods of protein synthesis, based on the ascertained amino acid sequences. In general, these methods employ the sequential addition of one or more amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid is protected by a suitable protecting group. The protected or derivatized amino acid can then be either attached to an inert solid support or utilized in

solution by adding the next amino acid in the sequence having the complementary (amino or carboxyl) group suitably protected, under conditions that allow for the formation of an amide linkage. The protecting group is then removed from the newly added amino acid residue and the next amino acid (suitably protected) is then added, and so forth. After the desired amino acids have been linked in the proper sequence, any remaining protecting groups (and any solid support, if solid phase synthesis techniques are used) are removed sequentially or concurrently, to render the final polypeptide. By simple modification of this general procedure, it is possible to add more than one amino acid at a time to a growing chain, for example, by coupling (under conditions which do not racemize chiral centers) a protected tripeptide with a properly protected dipeptide to form, after deprotection, a pentapeptide. See, e.g., J. M. Stewart and J. D. Young, *Solid Phase Peptide Synthesis*, 2nd Ed., Pierce Chemical Co., Rockford, IL (1984) and G. Barany and R. B. Merrifield, *The Peptides: Analysis, Synthesis, Biology*, editors E. Gross and J. Meienhofer, Vol. 2, Academic Press, New York, (1980), pp. 3-254, for solid phase peptide synthesis techniques; and M. Bodansky, *Principles of Peptide Synthesis*, Springer-Verlag, Berlin (1984) and E. Gross and J. Meienhofer, Eds., *The Peptides: Analysis, Synthesis, Biology*, *supra*, Vol. 1, for classical solution synthesis.

Recombinant human monoclonal antibody specific binding molecules can be prepared from the Fab molecules using known techniques. Bender et al. (1992) *Hum Antibod Hybridomas* 4:74. In particular, the coding sequence for the heavy chain portion of a selected Fab clone can be inserted into an expression vector along with the coding sequence for the constant domains of a human Ig heavy chain, using the various recombinant techniques described above. For example,

the mammalian expression vector pSG5 (Green et al. (1988) *Nucleic Acids Res* 16:369) can be used for this purpose.

Cloning involves overlap PCR to remove the
5 bacterial leader sequence (from the phagemid vector)
and to modify the N-terminus of the heavy chain coding
sequence to a human consensus sequence. The coding
sequence for the light chain portion of the selected
Fab clone can likewise be N-terminal modified to
10 include a human consensus sequence, and cloned into an
expression vector such as PSG5. The PSG5 vectors
contain an SV40 origin of replication such that, on
cotransfection of the heavy and light chain vectors
into mammalian cells, such as COS-7 cells, functional
15 antibody molecule production can be confirmed. Burton
et al. (1994) *Science* 266:1024-1027.

The heavy and light chains can subsequently
be cloned into separate cloning vectors, and either
the heavy or the light chain coding sequence subcloned
20 into the other vector to provide a combinatorial
plasmid. For example, the heavy and light chain
coding sequences can be respectively inserted into
pEE6 and pEE12 vectors (Bebbington et al. (1992)
Bio/Technology 10:169) which include a human
25 cytomegalovirus promoter and the glutamine synthetase
selectable marker. The heavy chain, along with
control elements from the PEE6 vector can then be
subcloned into the PEE12 vector to provide a
combinatorial plasmid. The combinatorial plasmid can
30 be expressed in a CHO cell expression system. Those
clones from the CHO expression system which provide
sufficient levels of recombinant antibody production
can be selected for scale-up. The recombinant
antibodies expressed in the CHO-system can be purified
35 using known techniques (e.g., affinity chromatography
using protein A), and the binding affinity of the

recombinant specific binding molecules assessed using an ELISA inhibition assay as described above.

Alternatively, the coding sequences for the Fab clones can be transferred into the vectors pcLCHC and pcIgG1, respectively, and then expressed as whole IgG in CHO cells as previously described. Samuelsson et al. (1996) *Eur. J. Immunol.* 26:3029.

Recombinant F(ab')₂ and recombinant Fv specific binding molecules can also be prepared from the phage-derived Fab clones using known techniques. Fv molecules generally comprise a non-covalently bound heavy chain:light chain heterodimer which includes the antigen-binding portion of the Fab molecule and retains much of the antigen recognition and binding capabilities of native antibody molecules. Inbar et al. (1972) *Proc. Nat. Acad. Sci. USA* 69:2659-2662; Hochman et al. (1976) *Biochem* 15:2706-2710; and Ehrlich et al. (1980) *Biochem* 19:4091-4096. Typically, the above-noted recombinant techniques used to construct the recombinant monoclonal antibodies can be modified to provide the truncated specific binding molecules. These molecules can also be cloned into CHO expression systems, purified and characterized as above.

The phage-derived Fab clones can further be used to provide single chain Fv (Sfv) molecules using known techniques. These Sfv molecules comprise a covalently linked heavy chain:light chain heterodimer which is expressed from a gene fusion including the heavy and light chain coding sequences obtained from the phage-derived Fab molecule, wherein the chains are linked by a peptide-encoding linker. Huston et al. (1988) *Proc. Nat. Acad. Sci. USA* 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated--but chemically separated--heavy and light chains into an Sfv molecule which will fold into a

three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Patent Nos. 5,091,513 and 5,132,405, to Huston et al.; and 4,946,778, to Ladner et al.

5 In the practice of the invention, recombinant DNA design methods are used to develop appropriate chemical structures for linking the heavy and light chains into the Sfv binding molecule. Design criteria include determination of the
10 appropriate length to span the distance between the C-terminus of one chain and the N-terminus of the other, wherein the linker is generally formed from small hydrophilic amino acid residues that do not tend to coil or form secondary structures. Such methods have
15 been described in the art. See, e.g., U.S. Patent Nos. 5,091,513 and 5,132,405 to Huston et al.; and U.S. Patent No. 4,946,778 to Ladner et al.

 The first general step of linker design involves identification of plausible sites to be
20 linked. Appropriate linkage sites on each of the immunoglobulin chains include those which will result in the minimum loss of residues from the heavy and light chains, and which will necessitate a linker having a minimum number of residues consistent with
25 the need for molecule stability. A pair of sites defines a "gap" to be linked. Linkers connecting the C-terminus of one chain to the N-terminus of the next generally include hydrophilic amino acids which assume an unstructured configuration in physiological
30 solutions and preferably are free of residues having large side groups which might interfere with proper folding of the heavy and light chains. Thus, suitable linkers would include polypeptide chains of alternating sets of glycine and serine residues, and
35 may include glutamic acid and lysine residues inserted to enhance solubility. One particular linker used in the practice of the invention has the amino acid

sequence [(Gly),Ser],. Another particularly preferred linker has the amino acid sequence comprising 2 or 3 repeats of [(Ser),Gly], such as [(Ser),Gly],.

5 Nucleotide sequences encoding such linker moieties can be readily provided using various oligonucleotide synthesis techniques known in the art. See, e.g., Sambrook, and Maniatis, *supra*.

Once the appropriate linker sequence has been ascertained, nucleotide sequences encoding the
10 Sfv molecules can be joined using an overlap PCR approach. See, e.g., Horton et al. (1990) *BioTechniques* 8:528-535. The ends of the light and heavy chains that are to be joined through the selected linker sequence are first extended by PCR
15 amplification of each chain, using primers that contain the terminal sequence of the chain region followed by all or most of the desired linker sequence. After this extension step, the light and heavy chains contain overlapping extensions which
20 jointly contain the entire linker sequence, and which can be annealed at the overlap and extended by PCR to obtain the complete Sfv sequence using methods known in the art. Genes present in expression cassettes comprising the sFv sequence can then be expressed in a
25 suitable expression system, and the sFv molecules produced therefrom can be purified and characterized as described above.

Vaccine Compositions

30 Therapeutic and prophylactic vaccine compositions are provided herein, which generally comprise mixtures of one or more of the above-described anti-HCV monoclonal antibodies, including Fab molecules, Fv fragments, sFv molecules and
35 combinations thereof. The prophylactic vaccines can be used to prevent HCV infection, and the therapeutic vaccines used to treat individuals following HCV

infection. Prophylactic uses include the provision of increased antibody titer to HCV in a vaccinated subject. In this manner, subjects at high risk of contracting HCV infection (e.g., immunocompromised individuals, organ transplant patients, individuals obtaining blood or blood product transfusions, and individuals in close personal contact with HCV-infected individuals) can be provided with passive immunity to the HCV agent. Furthermore, due to the cross-reactivity of the monoclonal antibodies, Fab molecules and sFv molecules produced herein, a level of protection is afforded against a number of heterologous HCV isolates. Other prophylactic uses for the present anti-HCV vaccines includes prevention of HCV disease in an individual after exposure to the infectious agent. Therapeutic uses of the present vaccines involve both reduction and/or elimination of the infectious agent from infected individuals, as well as the reduction and/or elimination of circulating HCV and the possible spread of the disease.

The compositions can be administered in conjunction with ancillary immunoregulatory agents, for example, cytokines, lymphokines, and chemokines, including but not limited to IL-2, modified IL-2 (cys125→ser125), GM-CSF, IL-12, γ -interferon, IP-10, MIP1 β and RANTES. When the vaccine compositions are used as therapeutic vaccines, the compositions can be administered in conjunction with known anti-HCV therapeutics, such as α -interferon (α -IFN) therapy which generally entails administration of 3 million units of α -IFN three times a week subcutaneously (Causse et al. (1991) *Gastroenterology* 101:497-502, Davis et al. (1989) *N Engl J Med* 321:1501-1506, Marcellin et al. (1991) *Hepatology* 13:393-397), interferon β (β -IFN) therapy (Omata et al. (1991) *Lancet* 338:914-915), ribivirin therapy (Di Bisceglie

et al. (1992) *Hepatology* 16:649-654, Reichard et al. (1991) *Lancet* 337:1058-1061) and antisense therapy (Wakita et al. (1994) *J Biol Chem* 269:14205-14210). Therapeutic vaccine compositions comprising the
5 present monoclonal antibodies can also be used in conjunction with known anti-HCV combination therapies, for example, the combination of α -IFN and ursodiol (Bottelli et al. (1993) (Abstr.) *Gastroenterology* 104:879, O'Brien et al. (1993) (Abstr.)
10 *Gastroenterology* 104:966) and the combination of β -IFN and ribivirin (Kakumu et al. (1993) *Gastroenterology* 105:507-512).

The preparation of vaccine compositions containing one or more antibodies, antibody fragments,
15 sFv molecules or combinations thereof, as the active ingredient is generally known to those of skill in the art. Typically, such vaccines are prepared as injectables (e.g., either as liquid solutions or suspensions or as solid forms suitable for solution or
20 suspension in liquids prior to injection). The compositions will generally also include one or more "pharmaceutically acceptable excipients or vehicles" such as water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. Additionally,
25 minor amounts of auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. The vaccine compositions may be emulsified or the active ingredient (monoclonal antibodies) may be encapsulated
30 in liposomes.

Once formulated, the vaccine compositions are conventionally administered parenterally, e.g., by injection (either subcutaneously or intramuscularly). Additional formulations suitable for other modes of
35 administration include oral and pulmonary formulations, suppositories, and transdermal applications. For suppositories, traditional binders

and carriers may include, for example, polyalkylene glycols or triglycerides. Such suppository formulations may be provided from mixtures containing the active ingredient(s) in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. The vaccine compositions are administered to the subject to be treated in a manner compatible with the dosage formulation, and in an amount that will be prophylactically and/or therapeutically effective. The amount of the composition to be delivered, generally in the range of from 1 to 500 micrograms of active agent per dose, depends on the subject to be treated, the capacity of the subject's immune system to mount its own immune-responses, and the degree of protection desired. The exact amount necessary will vary depending on the age and general condition of the individual to be treated, the severity of the condition being treated and the particular anti-HCV agent selected and its mode of administration, among other factors. An appropriate effective amount can be readily determined by one of skill in the art. Thus, a "therapeutically effective amount" of the composition will be sufficient to bring about treatment or prevention of HCV disease symptoms, and will fall in a relatively broad range that can be determined through routine trials.

In addition, the vaccine compositions can be given in a single dose schedule, or preferably in a multiple dose schedule. A multiple dose schedule is one in which a primary course of vaccination may be with 1-10 separate doses, followed by other doses given at subsequent time intervals needed to maintain or reinforce the action of the compositions. Thus,

the dosage regimen will also, at least in part, be determined based on the particular needs of the subject to be treated and will be dependent upon the judgement of the reasonably skilled practitioner.

5

Gene Therapy

The recombinant monoclonal antibodies can also be used for gene therapy. In this regard, genes encoding the recombinant antibodies can be introduced
10 into a suitable mammalian host cell for expression or coexpression using a number of viral based systems which have been developed for gene transfer into mammalian cells. For example, retroviruses provide a convenient platform for gene delivery systems. A
15 selected nucleotide sequence encoding a V_H and/or a V_L domain polypeptide can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to a subject. A number of
20 suitable retroviral systems have been described (U.S. Patent No. 5,219,740; Miller and Rosman (1989) *BioTechniques* 7:980-990; Miller, A.D. (1990) *Human Gene Therapy* 1:5-14; Scarpa et al. (1991) *Virology* 180:849-852; Burns et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:8033-8037; and Boris-Lawrie and Temin
25 (1993) *Cur. Opin. Genet. Develop.* 3:102-109. Particularly preferred methods for producing and using retroviral vectors for gene therapy herein are described, for example, in International Publication
30 No. WO 91/02805, published 7 March 1991, and in U.S. Patent Application Serial Nos.: 08/404,796, filed March 15, 1995 for "Eukarotic Layered Vector Initiation Systems;" 08/405,627, filed March 15, 1995 for "Recombinant α -Viral Vectors;" and 08/156,789,
35 filed November 23, 1993 for "Packaging Cells."

A number of suitable adenovirus vectors have also been described. Unlike retroviruses which

integrate into the host genome, adenoviruses persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis (Haj-Ahmad and Graham (1986) *J. Virol.* 57:267-274; Bett et al. (1993) *J. Virol.* 67:5911-5921; Mittereder et al. (1994) *Human Gene Therapy* 5:717-729; Seth et al. (1994) *J. Virol.* 68:933-940; Barr et al. (1994) *Gene Therapy* 1:51-58; Berkner, K.L. (1988) *BioTechniques* 6:616-629; and Rich et al. (1993) *Human Gene Therapy* 4:461-476).

10 Various adeno-associated virus (AAV) vector systems have been developed recently for gene delivery. Such systems can include control sequences, such as promoter and polyadenylation sites, as well as selectable markers or reporter genes, enhancer

15 sequences, and other control elements which allow for the induction of transcription. AAV vectors can be readily constructed using techniques well known in the art. See, e.g., U.S. Patent Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070

20 (published 23 January 1992) and WO 93/03769 (published 4 March 1993); Lebkowski et al. (1988) *Molec. Cell. Biol.* 8:3988-3996; Vincent et al. (1990) *Vaccines 90* (Cold Spring Harbor Laboratory Press); Carter, B.J. (1992) *Current Opinion in Biotechnology* 3:533-539;

25 Muzyczka, N. (1992) *Current Topics in Microbiol. and Immunol.* 158:97-129; Kotin, R.M. (1994) *Human Gene Therapy* 5:793-801; Shelling and Smith (1994) *Gene Therapy* 1:165-169; and Zhou et al. (1994) *J. Exp. Med.* 179:1867-1875.

30 Additional viral vectors which will find use for delivering the present nucleic acid molecules encoding the Fab molecules include those derived from the pox family of viruses, including vaccinia virus and avian poxvirus. By way of example, vaccinia virus

35 recombinants expressing the genes can be constructed as follows. The DNA encoding the particular Fab molecule is first inserted into an appropriate vector

so that it is adjacent to a vaccinia promoter and flanking vaccinia DNA sequences, such as the sequence encoding thymidine kinase (TK). This vector is then used to transfect cells which are simultaneously
5 infected with vaccinia. Homologous recombination serves to insert the vaccinia promoter plus the gene encoding the Fab molecule into the viral genome. The resulting TK recombinant can be selected by culturing the cells in the presence of 5-bromodeoxyuridine and
10 picking viral plaques resistant thereto.

A vaccinia based infection/transfection system can be conveniently used to provide for inducible, transient expression of the Fab molecules in a host cell. In this system, cells are first
15 infected in vitro with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the
20 polynucleotide of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into protein by the host translational machinery. The method provides
25 for high level, transient, cytoplasmic production of large quantities of RNA and its translation products. See, e.g., Elroy-Stein and Moss, *Proc. Natl. Acad. Sci. USA* (1990) 87:6743-6747; Fuerst et al., *Proc. Natl. Acad. Sci. USA* (1986) 83:8122-8126.

30 Alternatively, avipoxviruses, such as the fowlpox and canarypox viruses, can also be used to deliver the Fab-encoding nucleotide sequences. The use of an avipox vector is particularly desirable in human and other mammalian species since members of the
35 avipox genus can only productively replicate in susceptible avian species and therefore are not infective in mammalian cells. Methods for producing

recombinant avipoxviruses are known in the art and employ genetic recombination, as described above with respect to the production of vaccinia viruses. See, e.g., the International Publications WO 91/12882; WO 89/03429, published 20 April 1989; and WO 92/03545, published 5 March 1992.

Molecular conjugate vectors, such as the adenovirus chimeric vectors described in Michael et al. *J. Biol. Chem.* (1993) 268:6866-6869 and Wagner et al. *Proc. Natl. Acad. Sci. USA* (1992) 89:6099-6103, can also be used for gene delivery under the invention.

Assay Reagents and Diagnostic Kits

The above-described anti-HCV binding molecules (the recombinant monoclonal antibodies, including Fab molecules, Fv fragments and sFv molecules) which are capable of reacting immunologically with samples containing HCV particles are also used herein to detect the presence of HCV viral particles and/or viral antigens in specific binding assays of biological samples. In particular, the novel specific binding molecules of the present invention can be used in highly sensitive methods for screening and identifying individuals carrying and/or infected with HCV, as well as for screening for HCV-contaminated blood or blood products. The present binding molecules can also be used in assays for monitoring the progress of anti-HCV therapies in treated individuals, and for monitoring the growth rate of HCV cultures used in research and investigation of the HCV agent.

The format of specific binding assays will be subject to a great deal of variation in accordance with procedures that are well known in the art. For example, specific binding assays can be formatted to utilize one, or a mixture of several, of the

recombinant human monoclonal antibodies, (including Fab molecules, Fv fragments as well as sFv molecules) that have been prepared according to the present invention. The assay format can be generally based, for example, upon competition, direct binding reaction or sandwich-type assay techniques. Furthermore, the present assays can be conducted using immunoprecipitation or other techniques to separate assay reagents during, or after commencement of, the assay. Other assays can be conducted using specific binding molecules that have been insolubilized prior to commencement of the assay. In this regard, a number of insolubilization techniques are well known in the art, including, without limitation, insolubilization by adsorption to an immunoadsorbant or the like, absorption by contact with the wall of a reaction vessel, covalent crosslinking to insoluble matrices or "solid phase" substrates, noncovalent attachment to solid phase substrates using ionic or hydrophobic interactions, or by aggregation using precipitants such as polyethylene glycol or cross-linking agents such as glutaraldehyde.

There are a large number of solid phase substrates which can be selected for use in the present assays by those skilled in the art. For example, latex particles, microparticles, magnetic-, para-magnetic- or nonmagnetic-beads, membranes, plastic tubes, walls of microtitre wells, glass or silicon particles and sheep red blood cells all are suitable for use herein.

In general, most of the present assays involve the use of a labeled binding complex formed from the combination of a specific binding molecule (recombinant monoclonal antibodies, Fab fragments, Fv fragments and sFv molecules) with a detectable label moiety. A number of such labels are known in the art and can be readily attached (either using covalent or

non-covalent association techniques) to the binding molecules of the present invention to provide a binding complex for use in the above-noted assay formats. Suitable detectable moieties include, but
5 are not limited to, radioactive isotopes, fluorescers, luminescent compounds (e.g., fluorescein and rhodamine), chemiluminescers (e.g., acridinium, phenanthridinium and dioxetane compounds), enzymes (e.g., alkaline phosphatase, horseradish peroxidase
10 and beta-galactosidase), enzyme substrates, enzyme cofactors, enzyme inhibitors, dyes, and metal ions. These labels can be associated with the binding molecules using attachment techniques that are known in the art.

15 Exemplary assay methods generally involve the steps of: (1) preparing the detectably labeled binding complexes as above; (2) obtaining a sample suspected of containing HCV particles and/or HCV antigen; (3) incubating the sample with the labeled
20 complexes under conditions which allow for the formation of a specific binding molecule-antigen complex (e.g., an antibody-antigen complex); and (4) detecting the presence or absence of labeled binding molecule-antigen complexes. As will be appreciated by
25 those skilled in the art upon the reading of this specification, such assays can be used to screen for the presence of HCV infection in human donor blood and serum products, for monitoring the growth rate of HCV cultures in diagnostic and/or research settings, for
30 detecting HCV infection in an individual, or for monitoring the therapeutic effect of an anti-HCV treatment protocol in an infected subject. When the assays are used in the clinical setting, e.g., for detecting HCV infection or monitoring anti-HCV
35 therapies, samples can be obtained from human and animal body fluids, such as whole blood, serum, plasma, cerebrospinal fluid, urine and the like.

Furthermore, the assays can be readily used to provide quantitative information using reference to standards or calibrants as known in the art.

5 In one particular assay method of the invention, an enzyme-linked immunosorbent assay (ELISA) can be used to quantify an HCV antigen concentration in a sample. In the method, the specific binding molecules of the present invention are conjugated to an enzyme to provide a labeled
10 binding complex, wherein the assay uses the bound enzyme as a quantitative label. In order to measure antigen, a binding molecule capable of specifically binding the selected HCV antigen (e.g., an antibody molecule) is immobilized to a solid phase substrate
15 (e.g., a microtitre plate or plastic cup), incubated with test sample dilutions, washed and incubated with the binding molecule-enzyme complexes of the invention, and then washed again. In this regard, suitable enzyme labels are generally known, including, for example, horseradish peroxidase. Enzyme activity
20 bound to the solid phase is measured by adding the specific enzyme substrate, and determining product formation or substrate utilization colorimetrically. The enzyme activity bound to the solid phase substrate
25 is a direct function of the amount of antigen present in the sample.

In another particular assay method of the invention, the presence of HCV in a biological sample (e.g., as an indicator of HCV infection) can be
30 detected using strip immunoblot assay (SIA) techniques, such as those known in the art which combine traditional Western and dot blotting techniques, e.g., the RIBA® (Chiron Corp., Emeryville, CA) test. In these assays, one or more of the
35 specific binding molecules (the recombinant monoclonal antibodies, including Fab molecules) are immobilized as individual, discrete bands on a membranous support

test strip. Visualization of reactivity with HCV particles present in the biological sample is accomplished using sandwich binding techniques with labeled antibody-conjugates in conjunction with a colorimetric enzyme substrate. Internal controls can also be present on the strip. The assay can be performed manually or used in an automated format.

Furthermore, the recombinant human monoclonal antibodies, (including Fab molecules, Fv fragments as well as sFv molecules) that have been prepared according to the present invention can be used in affinity chromatography techniques in order to detect the presence of HCV in a biological sample. Such methods are well known in the art.

Kits suitable for use in conducting any of the above-described assays and affinity chromatography techniques, and containing appropriate labeled binding molecule complex reagents can also be provided in accordance with the practice of the invention. Assay kits are assembled by packaging the appropriate materials, including all reagents and materials necessary for conducting the assay in a suitable container, along with an appropriate set of assay instructions.

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

35

ExperimentalExample 1Characterization of the Library Donor

5 Bone marrow was obtained from a 60 year old
asymptomatic, male blood donor, who was found to be
HCV positive during regular screening in conjunction
with a blood donation. The cause of infection was
unknown. The donor was unimmunized, and had received
10 no treatment for the HCV infection prior to the bone
marrow aspiration, which amounted to approximately 3
ml. The genotype of HCV in the donor's serum at the
time of bone marrow donation was determined using a
published method and found to be HCV 2b. Widell et
15 al. (1994) *J Med Virol* 44:272-279.

In particular, the nucleotide sequence of
the hypervariable region 1 (HVR1) of the E2 gene of
the HCV isolate from the donor was obtained as
follows. Single stranded template DNA was obtained,
20 and a sequencing reaction was performed using newly
designed primers (A. Widell et al, manuscript in
preparation), in a cycle sequencing reaction with
labelled nucleotides (PCR cycle sequencing kit,
Perkin-Elmer) according to the manufacturers
25 instructions. Allander et al. (1994) *J Med Virol*
43:415-419. The reaction product was run on an
automated sequencer (Applied Biosystems, CA), and the
data edited and analyzed using MacMolly software
(available from SoftGene, Berlin, Germany). The
30 deduced amino acid sequence of the HVR-E2 region of
the donor isolate was determined to be as follows:
VAGVDASTYTTGGQSGRTTYGIVGLFSLGPSQKLSLINTNGSWHINR (SEQ
ID NO:).

35

Example 2Construction of the Phage Display Library

Lymphocytes were isolated from the bone marrow sample obtained in Example 1 using Ficoll-Paque (Pharmacia Biotech, Uppsala, Sweden). Total RNA was extracted by the acid phenol extraction method (Chomczynski et al. (1987) *Anal Biochem* 162:156-159), first strand cDNA synthesis utilizing oligo-dT priming of 10 µg of RNA was performed (cDNA synthesis kit, Pharmacia Biotech) and heavy (Fd) and light chain DNA was PCR amplified using 5' biotinylated primers of previously published sequences for γ1 heavy and κ light chains (5' primers: VH1a, VH1f, VH2f, VH3a, VH3f, VH4f, VH6a, VH6f, Vk1a, Vk2a, Vk3a; 3'-primers: CG1z and CK1a) (available from Scandinavian Gene Synthesis, Köping, Sweden). See, e.g., Persson et al. (1991) *Proc Natl Acad Sci USA* 88:2432-2436, Kang et al. (1991) *Methods: Comp. Meth Enzymol* 2:111-118.

PCR was performed using a Thermal Cycler 4800 (Perkin-Elmer) at 94°C for 5 minutes, then 35 cycles of 94° for 60 seconds, 52° for 30 seconds, and 72° for 180 seconds. After the 35 cycles, an extension step at 72° for 10 minutes completed the PCR procedure. PCR products were analyzed by running a fraction of each in a 1.5% agarose gel, the heavy and light chain DNA were pooled separately, phenol-chloroform extracted and then ethanol precipitated.

12 µg each of heavy and light chain DNA was gel purified on a 2.5% agarose gel, electroeluted (Schleicher & Schuell, Germany), and digested with the restriction endonucleases *XhoI/SpeI*, and *SacI/XbaI*, respectively (available from Life Technologies, Gaithersburg, MD). The digested PCR products were subsequently gel purified, recovered by electroelution, and ligated into the vector pComb3H (Barbas, III et al. (1995) *Methods: Comp. Meth Enzymol* 8:94-103) after it had been digested with the

corresponding restriction enzymes and gel purified/electroeluted as previously described for similar vector systems (Yang et al. (1995) *J Mol Biol* 254:392-403, Barbas, III et al. (1991) *Proc Natl Acad Sci USA* 88:7978-7982). For ligations, T4 ligase (Life Technologies) was used at 0.5 units per 10 μ l reaction volume at approximately 16°C over night. The combinatorial library was expressed on phage, including harvesting of phage, as reported.

10 Samuelsson et al. (1995) *Virology* 207:495-502.

Ligation with light chain genes into pComb3H gave a library of 2 x 10⁷ members. The subsequent ligation of Fd genes into this library resulted in a γ l/ κ library with 2 x 10⁶ members.

15

Example 3

Preparation of the HCV E2 Selecting Antigens

A truncated, secreted form of the HCV E2 molecule was constructed which includes amino acids 383_{ala} through 715_{lys} (using the nomenclature of Choo et al. (1991) *Proc Natl Acad Sci USA* 88:2451-2455). The E2 molecule was expressed using a Chinese hamster ovary cell/dihydrofolate reductase (CHO/DHFR) expression system to provide a "conformational HCV E2 antigen" as follows. A DNA fragment of HCV E2 from amino acid 383 to amino acid 715 of HCV1 was generated by PCR and then ligated into a plasmid vector having the murine cytomegalovirus (MCMV) immediate early promoter/enhancer (Dorsch-Hasler et al. (1985) *Proc Natl Acad Sci USA* 82:8325-8329) and the selectable *dhfr* gene marker. The resultant plasmid was then stably transfected into *dhfr* CHO cells to generate a stable recombinant CHO cell line which secreted the conformational HCV E2 antigen.

35 The conformational E2 antigen was purified using known methods (Rosa et al. (1996) *Proc Natl Acad Sci USA* 93:1759-1763) as follows. Conditioned media

from the CHO cells was concentrated 15-fold by ultrafiltration, followed by a further 10-fold volume reduction by ammonium sulfate precipitation at 75% saturation, and redissolution into 25 mM Tris chloride/1 mM EDTA, pH 7.5. The monoclonal antibody 5E5/H7 (raised against HeLa E1/E2) was used for purification. The antibody column was equilibrated in 25 mM Tris chloride/0.15 M NaCl, pH 7.5. The ammonium sulfate-precipitated E2 was dissolved in 25 mM Tris chloride/1 mM EDTA, pH 7.5, and loaded onto the column. The column was washed with phosphate buffered saline (PBS)/1 M NaCl and then eluted with 3-4 column volumes of Actisep (Sterogene, Arcadia, CA). All of the yellow-colored Actisep-containing fractions were pooled, concentrated in a stirred cell ultrafilter, and diafiltered into PBS buffer.

A recombinant HCV E1/E2 complex antigen was constructed and expressed using a Chinese hamster ovary cell/dihydrofolate reductase (CHO/DHFR) expression system as follows. Plasmid pMCMV-HC5p (Spaete et al. (1992) *Virology* 188:819-830) which encodes the HCV structural region as a 917 amino acid polypeptide spanning Met₁ to Gly₉₁₇ of the HCV1 genome was generated by cloning a 2813 base pair (bp) HCV *StuI* fragment from pGEM-4 blue-HC5p-1 8 into the unique *SalI* site of the mammalian cell expression vector pMCVAdhfr (Spaete et al. (1990) *J Virol* 64:2922-2931). The pMCVAdhfr vector encodes the selectable *dhfr* gene with transcription of the expressed gene driven by the MCMV immediate early promoter/enhancer and terminated by SV40 polyadenylation sequences. The Klenow fragment (Boehringer-Mannheim Biochemicals, Indianapolis, IN) was used to fill the *SalI* site prior to ligation of the *StuI* fragment. The pMCMV-HC5p plasmid was stably transfected into *dhfr* CHO cells to generate a stable

recombinant CHO cell line (#62) which expresses the recombinant HCV E1/E2 complex antigen.

The recombinant E1/E2 complex antigen was purified using known methods (Choo et al. (1994) *Proc Natl Acad Sci USA* 91:1294-1298) as follows. CHO cell line #62 was harvested by pelleting and freezing. After lysis by Dounce homogenization in hypotonic buffer, the pellet was extracted by homogenizing in 2% Triton X-100®, and E1 (33 kD) and E2 (72 kD) were selectively copurified by successive chromatography on agarose-bound Galanthus Nivalis-lectin (Vector Laboratories) and fast-flow S-Sepharose cation exchanger (Pharmacia).

15

Example 4

Panning of the Combinatorial Library

Antigenic selection of specifically-binding Fab molecules was conducted using a variation on known techniques. Burton et al. (1991) *Proc Natl Acad Sci USA* 88:10134-10137. In particular, four wells of a microtiter plate (Costar 3690, Cambridge, MA) were coated at 4°C overnight with 50 µl of either purified recombinant conformational HCV E2 antigen or purified recombinant E1/E2 complex antigen (prepared in Example 3), expressed in CHO cells, at 2.5 µg/ml. Spaete et al. (1992) *Virology* 188:819-830. Blocking was effected by completely filling the wells with 5% non-fat dry milk in PBS for 1 h at 22°C. 50 µl of the phage library (5×10^{10} cfu) was added to each well, and the plate was incubated for 2 hours at 37°C. Phage were removed and each well was washed by completely filling with a solution of PBS and 0.5% Tween 20 for 5 minutes, and then thoroughly removing the wash solution. Washing was performed 1-10 times as described below. Phage were eluted by adding 50 µl per well of elution buffer (0.1M HCl adjusted to pH 2.2 with solid glycine) and incubating for 10 minutes at

ambient temperature. The elution buffer was removed and neutralized with 3 μ l of 2M Tris base per 50 μ l of elution buffer. *E. coli* XL-1 blue cells (Barbas, III et al. (1991) *Methods: Comp. Meth Enzymol* 2:119-124) were infected by the eluted phage, aliquots plated, and propagation of phage after each round of panning effected as has been described. Samuelsson et al. (1995) *Virology* 207:495-502.

In the first group of pannings (Panning Series I), the number of washings was increased in each subsequent panning round for three rounds (1, 3 and 10 washings, respectively), while in the second group of pannings (Panning Series II), a single panning round with 10 washings was performed. As can be seen by the results depicted in Table I, a 100 fold increase in eluted phage was noted in Panning Series I.

20

TABLE I			
Panning No.	Washes (No.)	Eluted Phage (cfu)	Enrichment Factor
<i>Panning Series I</i>			
1	1	2.6×10^6	--
2	3	2.3×10^7	9
3	10	2.6×10^8	100
<i>Panning Series II</i>			
1	10	2.4×10^5	

25

The antigen used in both the Series I and II pannings was the recombinant conformational E2 antigen. A third group of pannings (Panning Series III) was performed as in Series I; however, the recombinant E1/E2 complex antigen was used to select positive clones.

35

Example 5

Expression of the Fab Molecules

Fab molecules were expressed by growing ampicillin resistant *E. coli* XL-1 blue cell colonies, containing Fab plasmids, with the *gIII* gene (encoding the cpIII anchor protein) (a) intact to provide insoluble Fab fragments, or (b) deleted by digestion with *SpeI* and *NheI* to provide soluble Fab fragments (digestion with these enzymes provides compatible cohesive ends; thus, the resulting DNA fragment lacking the *gIII* fragment can be gel-purified and self-ligated), in SB medium (super broth; 30 g tryptone, 20 g of yeast extract, and 10 g of MOPS per liter, pH 7) (Burton et al. (1991) *Proc Natl Acad Sci USA* 88:10134-10137) containing 50 µg/ml ampicillin and 1 % glucose, until an OD_{600nm} of about 1.0 was reached. The bacterial host cells were pelleted by centrifugation, and media exchanged to SB medium with 1 mM IPTG and 20 mM MgCl₂, and the cells resuspended. The resulting culture was incubated at room temperature on a shaker platform set at about 290 rpm and left overnight. The following day, cells were spun down, the supernatant discarded, PBS added (to between 2 to 4% of original culture volume) and the periplasmic contents of the bacterial cells released by three cycles of freeze- thawing. Bacterial debris was pelleted by centrifugation, and the Fab molecule-containing supernatant aliquoted to new vials. The Fab molecules were maintained at -20°C until used.

Example 6

Expression Levels of the Fab Clones

The expression levels of Fab molecules (obtained in Example 5) were ascertained using known ELISA techniques. Samuelsson et al. (1995) *Virology* 207:495-502. In particular, goat anti-human F(ab'), (Pierce, USA) or goat anti-human Fd (The Binding Site,

UK) was diluted 1:1000 in 0.1 M carbonate-bicarbonate buffer, pH 9.6 and coated on microtiter wells by incubation overnight at 4°C. Coating solution was discarded, and the wells were blocked with 5% dry milk in PBS for 1 hr at ambient temperature, after which the blocking solution was removed, and Fab samples (from Example 5) at appropriate dilutions in PBS-T were added. After incubation at ambient temperature for 1 hour, the plates were washed, and ALP-goat anti human F(ab')₂ at a 1:500 dilution was added. After 1 hour incubation, and five subsequent washes, the label substrate solution, p-nitrophenylphosphate (Sigma, USA) in 0.1 M diethanolamine, pH 9.8 was added. Absorbance was measured at 405 nm in a microplate reader (Dynastar, MA). Most Fab clones were found to produce between 0.2 and 2.0 mg Fab/ L culture, corresponding to 10-100 µg/ml in the periplasmic preparations. The expressed Fab clones were screened for E2 reactivity and promptly sequenced, in order to identify multiple copies of the same original clone.

Example 7

Western Blot for Heavy and Light Chain Expression

In order to test for correct expression of both chains, several Fab molecules were analyzed in Western blots using antiserum for human Fd- and light chains. In particular, 10 µl of the periplasmic Fab molecule preparations (prepared in Example 5) were separated on a precast 12% Tris-glycine polyacrylamide gel, and transferred to a nitrocellulose membrane by electroblotting using an Xcell Mini-cell apparatus (Novex Experimental Technology, San Diego, CA). The membrane was blocked in 5% dry milk over night, and incubated with either alkaline phosphatase coupled anti-human Fab antiserum (available from Pierce) diluted 1:1000 in 5% dry milk and 0.05% Tween 20 (PBS-

MT) for 3 h at 22°C during constant rocking. The subject anti-human Fab antiserum was chosen since it is known to be mainly reactive to light chains. To detect the heavy (Fd) chain expression products, strips were first incubated with a sheep-anti-human Fd serum (Binding Site, U.K.) which was diluted to 1:1000 and incubated (as above), washed and then incubated again with a secondary antibody, AP-anti-goat IgG (Sigma, St. Louis, MO) at a dilution of 1:500.

Following the last incubation for 1 hour at 22°C, the membranes were washed three times in PBS-T, and color development was performed with 2 ml BCIP/NBT solution (Sigma, St Louis, MO) for 7 minutes. Membranes were rinsed in water and dried. Prestained molecular weight markers (Amersham, U.K.) were used in each blot.

For all clones tested, expression of both chains was approximately equivalent. The heavy chain (expressed as a fusion polypeptide with the truncated gIII protein) showed an approximate molecular weight of 70 kD.

Example 8

Sequencing of the Fab Clones

Plasmid DNA from each Fab molecule clone grown in the *E. coli* XL-1 blue cell cultures (in Example 5), was isolated using a Wizard mini prep DNA purification reagent system (Promega). Single stranded DNA was obtained by PCR, using primers that hybridized upstream and downstream of the cloning regions in the pComb3H vector (pC3H-2488S: 5'-CAA CGC AAT TAA TGT GAG TTA G (SEQ ID NO:); G-back: 5'-GCC CCC TTA TTA GCG TTT GCC ATC (SEQ ID NO:)). In each reaction, one of the two PCR primers used was biotinylated at the 5' terminus. After 35 cycles of PCR amplification, single stranded DNA was obtained by

denaturing the DNA under alkaline conditions and absorbing the biotinylated DNA strand to streptavidin coated beads (Dynal, Oslo, Norway) using known techniques. Hultman et al. (1989) *Nucl Acid Res* 17:4937-4945.

Dideoxy sequencing reactions according to the method of Sanger et al. (1977) *Proc Natl Acad Sci USA* 74:5463-5467 was performed utilizing FITC-labelled primers hybridizing 3' of the junction between the variable and constant Ig regions or 5' to the start of the heavy and light chain genes. Particularly, SEQKb: 5'-ATA GAA GTT GTT CAG CAG GCA (SEQ ID NO:) and omp-seq: 5'-AAG ACA GCT ATC GCG ATT GCA G (SEQ ID NO:) were used for the κ light chains. SEQGb: 5'-GTC GTT GAC CAG GCA GCC CAG (SEQ ID NO:) and pel-seq: 5'-ACC TAT TGC CTA CGG CAG CCG (SEQ ID NO:) were used for the γ heavy chains. The reaction products were run on an automated sequencer (A.L.F., Pharmacia Biotech), and were translated and aligned using the MacMolly software (SoftGene, Berlin, Germany).

From the first series of pannings conducted in Example 4 (Panning Series I), 10 Fab molecule clones (identified as Fab molecule clones L1-L10) that were assayed for expression (as described in Examples 6 and 7) were sequenced using the above-described sequencing method. These 10 clones were found to have very similar CDR3 sequences in their heavy chains (the H3 region), indicating that they all derived from the same B-cell clone. Litwin et al. (1990) *J Exp Med* 171:293-297. However, while the VDJ junctions and the length of the H3 regions were identical, a number of different point mutations were identified in their heavy chains, and each heavy chain was combined with a different light chain. Two clones, identified as Fab molecule clones L1 and L3, were selected for further testing.

From the second series of pannings conducted in Example 4 (Panning Series II), 20 Fab molecule clones (identified as Fab molecule clones 1:1-1:20) that were assayed for expression (as described in Examples 6 and 7) were sequenced. From this round of sequencing, 6 Fab clones were found to produce insufficient levels of Fab, and 4 Fab clones were found to exhibit cross reactivity to control antigen in a specific binding assay. From the sequencing information obtained from the remaining 10 Fab clones, it was found that 7 Fab clones carried heavy chains related to the ones found in the Panning Series I (the L1-10 Fab molecule clones). However, 3 of the 7 Fab clones had distinctly different H3 regions from the Panning Series I clones, and were also unique relative to each other. These clones were selected for further testing and identified herein as Fab molecule clones 1:5, 1:7, 1:11.

From the third series of pannings conducted in Example 4 (Panning Series III), 30 Fab molecule clones were assayed for expression (as described in Examples 6 and 7), and 16 were found to be reactive to both the E1/E2 complex antigen and to the E2 antigen alone. These 16 clones were sequenced as above. 12 of the 16 clones that were sequenced were found to have a H3 sequence similar to clones L1-L10, while the remaining 4 were found to have unique H3 sequences. Two of the 4 clones having unique H3 sequences were selected for further testing and are identified herein as Fab molecule clones A8 and A12.

The κ light chain nucleic acid sequences of the following Fab molecule clones: 1:5 (SEQ ID NO:); 1:7 (SEQ ID NO:); 1:11 (SEQ ID NO:); L3 (SEQ ID NO:); L1 (SEQ ID NO:); A8 (SEQ ID NO:); and A12 (SEQ ID NO:) are depicted in Figures 3A-3G, respectively.

The γ 1 heavy chain nucleic acid sequences of the following Fab molecule clones: 1:5 (SEQ ID NO:); 1:7 (SEQ ID NO:); 1:11 (SEQ ID NO:); L3 (SEQ ID NO:); L1 (SEQ ID NO:); A8 (SEQ ID NO:); and A12 (SEQ ID NO:) are depicted in Figures 4A-4G, respectively.

The deduced γ 1 heavy chain amino acid sequences of Fab molecule clones 1:5 (SEQ ID NO:); 1:7 (SEQ ID NO:); 1:11 (SEQ ID NO:); L3 (SEQ ID NO:); L1 (SEQ ID NO:); A8 (SEQ ID NO:); and A12 (SEQ ID NO:) are depicted in Figures 1A-1G, respectively. The CDR regions (CDR1, CDR2 and CDR3) from each chain have been identified in the Figures.

The deduced κ light chain amino acid sequences of Fab molecule clones 1:5 (SEQ ID NO:); 1:7 (SEQ ID NO:); 1:11 (SEQ ID NO:); L3 (SEQ ID NO:); L1 (SEQ ID NO:); A8 (SEQ ID NO:); and A12 (SEQ ID NO:) are depicted in Figures 2A-2G, respectively. The CDR regions (CDR1, CDR2 and CDR3) from each chain have also been identified as noted above.

In summary, out of 50 clones that were obtained from the three panning series, 36 were found to be specific to E2, and 29 of those 36 E2-specific clones share a related heavy chain.

Example 9

ELISA Assay for HCV E2 Antigen Reactivity

The Fab molecule clones 1:5, 1:7, 1:11 and L3 were screened for HCV E2 antigen reactivity as follows. Either recombinant conformational HCV E2 antigen, or recombinant HCV E1/E2 complex antigen (prepared as described in Example 3) was diluted to 0.25 μ g/ml in 0.05 M carbonate-bicarbonate buffer, pH 9.6, and coated to microtitre wells (Costar #3690; Life Technologies) overnight at 4°C. Unbound antigen was discarded, and the wells were blocked with 5%

nonfat dry milk in PBS for 60 minutes at ambient temperature. After the blocking solution was discarded, solutions containing the Fab molecules to be tested were added at 1:2, 1:10 and 1:100 dilutions (diluent: PBS with 0.1 % NP-40). The plates were incubated at ambient temperature for 2 hours, washed five times with PBS with 0.05% Tween 20 (PBS-T), and ALP-goat anti-human F(ab')₂ (Pierce, Rocherford, IL) was added at a 1:1000 dilution. After 60 minutes and subsequent washes, substrate solution (p-nitrophenylphosphate) (SIGMA, St. Louis, MO) was added and absorbance was measured at 405nm in a microplate reader (Dynastar, MA).

The cut-off value for positive readings was set at 4 times the OD value obtained for a negative control sample which comprised an anti-HIV Fab of equal concentrations. Barbas III et al. (1991) *Proc Natl Acad Sci USA* 88:7978-7982. For control purposes, bovine serum albumin (BSA) (SIGMA), HIV gp120_{LAI} (Intracell, Cambridge, MA) and tetanus toxoid (TT) (SBL Vaccin, Solna, Sweden) coated at 5, 1, and 1 µg/ml, respectively, were used in corresponding ELISAs as controls for unspecific reactivity.

The results from the ELISAs are depicted in Table II below. As can be seen, Fab molecules expressed from the 1:5, 1:7, 1:11 and L3 clones each reacted strongly with both the conformational HCV E2 antigen and the HCV E1/E2 complex antigen, while showing no cross reaction with the control antigens (BSA, HIVgp120 and TT).

35

TABLE II

Clone	[Fab] ($\mu\text{g/ml}$) ¹	ELISA reactivity to:				
		E2 ²	E1/E2 ²	BSA ²	HIV gp120 ²	TT ²
L 3	100	1.695	2.460	0.089	ND	0.313
1:5	10	0.219	0.614	0.031	ND	0.012 ³
1:7	100	>3.000	>3.000	0.006	ND	ND
1:11	10	>3.000	1.831	0.506	ND	ND

¹ Fab concentration in the periplasmic preparation used in the analyses.

² OD_{405 nm}, sample diluted 1:10.

³ sample diluted 1:100.

Example 10

Western Blot Assay for HCV E2 Antigen Reactivity

The Fab molecule clones 1:5, 1:7, 1:11 L1, L3, A8 and A12 were screened for HCV E2 antigen reactivity using the following techniques. Western blots were conducted using 1 μg of the recombinant conformational HCV E2 glycoprotein (obtained in Example 3) that was denatured by heating to 98°C for 5 min in a Laemmli buffer, separated on a 8-16% polyacrylamide gradient gel (Novex Experimental Technologies), and transferred to a nitro-cellulose membrane that was blocked as described above, and cut into strips. Subsequently, each strip was incubated with a Fab preparation (expressed from the 1:5, 1:7, 1:11, L3, L1, A8 and A12 clones) that was diluted 1:20 in PBS-MT for 2 hours at 22°C with constant rocking. The strips were washed three times in PBS-T, and alkaline phosphatase conjugated goat anti-human Fab serum (Pierce), diluted 1:1000 in PBS-MT, was added. Following incubation for 1 hour at 22°C, the strips were again washed three times in PBS with 0.05% Tween

20, and color development was performed with 2 ml
BCIP/NBT solution (Sigma, St Louis, MO) for 10
minutes. As a positive control, human anti-HCV
positive serum was incubated with one strip instead of
5 the Fab preparations.

None of the tested Fab molecules (from
clones 1:5, 1:7, 1:11, L3, L1, A8 and A12) reacted to
the denatured HCV E2 antigen in the Western blot,
indicating that each Fab molecule binds to a
10 conformational epitope of the HCV E2 antigen.
However, the positive control (human anti-HCV positive
serum) did react with the denatured E2 antigen in the
Western Blot.

The above-described assay was repeated under
15 identical conditions, with the single change being use
of HCV E2 antigen that was gel separated under non-
denaturing conditions. Both of the clones (1:7 and
A8) tested in this further assay were found to bind to
the non-denatured E2 antigen.

20

Example 11

Inhibition ELISA Assay for Affinity Determination

The affinity of the Fab molecules (from
clones 1:5, 1:7, 1:11, L3, L1, A8, and A12) for HCV E2
25 antigen was estimated using an inhibition ELISA method
as previously described. Persson et al. (1991) *Proc
Natl Acad Sci USA* 88:2432-2436, Rath et al. (1988) *J
Immun Methods* 106:245-249. Samples to be tested were
first titred at ten-fold dilutions in order to bracket
30 a concentration where a ten-fold reduced concentration
gave a substantial reduction in detected binding in
the HCV E2 ELISA. For affinity measurements, coating
of microtitre wells with HCV E2 antigen (HCV genotype
1a) and subsequent blocking was done as described
35 above for the ELISA conducted in Example 9.
Appropriate dilutions of the Fab samples, with or
without added soluble HCV E2 antigen (final

concentration 5 $\mu\text{g/ml}$) were added to the wells, and incubated at ambient temperature for 3 hours. The plates were washed uniformly 4 times with PBS-T, and developed using AP-anti Fab, substrate, and spectrophotometer reader as described above in Example 9. The reduction of OD in the presence of soluble HCV E2 antigen was calculated and the concentration needed for a 50% reduction estimated by extrapolation.

As depicted in Table III below, the approximate affinities of the Fab molecules (from clones 1:5, 1:7, 1:11, L3, L1, A8 and A12) for the recombinant conformational HCV E2 antigen (HCV genotype 1a), varied between 1×10^7 and $2 \times 10^8 \text{ M}^{-1}$.

15

20

25

TABLE III		
Clone	K_d^1 (nM)	Affinity (M^{-1})
L3	28	4×10^7
1:5	>100	$<1 \times 10^7$
1:7	6	2×10^8
1:11	28	4×10^7
L1	28	4×10^7
A8	6	2×10^8
A12	100	1×10^7

¹ Approximate concentration needed of soluble E2 for 50% reduction in OD.

The affinity of the Fab molecules (from clones 1:7, A8 and A12) for a different HCV E2 antigen (HCV genotype 1b) was also assessed using the above-described inhibition ELISA. The affinities for the E2 antigen of genotype 1b in each of the tested molecules was found to be similar to those reported above (Table III) for the genotype 1a E2 antigen.

In addition, whole recombinant IgG molecules prepared from the following Fab molecule clones: L1;

L3; 1:5; 1:7; and 1:11, were assessed using the above-described inhibition ELISA with the HCV genotype 1a E2 antigen. The affinities observed were similar to those reported above (Table III) for the Fab molecule clones.

Example 12

Inhibition of HCV E2 Binding

The ability of the Fab molecules (from clones 1:5, 1:7, 1:11, L3, L1, A8 and A12) to block the binding of HCV E2 to target cells was determined using the neutralization of binding (NOB) method of Rosa et al. (1996) *Proc Natl Acad Sci USA* 93:1759-1763. More particularly, purified conformational HCV E2 antigen (from both genotypes HCV 1a and HCV 1b, and prepared as described in Example 3) was used in indirect immunofluorescence experiments to assess the ability of two separate batches of bacterially expressed Fab molecule clones to neutralize binding of the HCV E2 polypeptide to human cells *in vitro*.

In the assay, 20 μ l of the purified conformational HCV E2 antigen (in PBS at 0.5 μ g/ml) was mixed with various dilutions of the Fab clones. After incubation at 4°C for 1 hour, pellets of MOLT-4 cells (a human cell line reported to allow low-level HCV replication *in vitro* as described by Shimizu et al. (1992) *Proc Natl Acad Sci USA* 89:5477-5481), were added and the resulting reaction mixture incubated for 1 hour at 4°C. Unbound HCV antigen and antibodies were removed by two centrifugations in PBS at 200 x g for 5 minutes at 4°C. The cells were then incubated for 30 minutes at 4°C with human anti-HCV E2 reactive serum. The cells were then washed twice in PBS and incubated for 30 minutes with fluorescein isothiocyanate-conjugated antiserum specific for human Fab. The cells were washed again in PBS at 4°C and resuspended in 100 μ l of PBS. Cell-bound fluorescence

was analysed with a flow cytometer (FACScan, Becton Dickinson) using Lysis II software (Becton Dickinson). Mean fluorescence intensity of cell populations incubated with the various Fab preparations were
5 calculated, and compared to mean fluorescence intensity of cells incubated without antibodies or without the E2 antigen.

The results are depicted below in Table IV. As can be seen, all seven of the tested Fab clones
10 efficiently inhibited MOLT-4 cell binding by the conformational HCV E2 antigen (both genotypes HCV 1a and HCV 1b). Clones A8, 1:7, L1 and L3 had very high neutralization activity in the assay. The 50% reduction titer is shown for all tested clones in
15 Table IV, and the complete assay result for 4 of the clones is shown in Figure 5. Two negative control Fab clones, prepared in the same manner as described above but directed to HIV-1 envelope glycoprotein gp120 (clones b12 and b14), did not have neutralization
20 activity in the assay. Fab clones expressed in eucaryotic cells, and recombinant whole IgG molecules derived from the Fab clones were found to be negative in a similar NOB assay.

25

30

35

Table IV			
	Antigen		
	E2 1a		E2 1b
	Exp. I	Exp. II	Exp. II
clone			
1:5	1.5*	2.5	2.5
1:7	0.02	0.01	0.02
1:11	0.4	0.1	0.15
L1	0.02	0.03	0.03
L3	0.02	0.02	0.03
A8	n.d.	0.001	0.007
A12	n.d.	0.1	0.15
b12	>10	>10	>10
b14	n.d.	>10	>10

* Fab concentration in $\mu\text{g/ml}$.

Since the first contact between the HCV virus and its host occurs via binding of the virus envelope to cell-surface receptors, the ability of the present Fab molecules to neutralize this interaction establishes the effectiveness of using those molecules in vaccinations to provide passive immunization to HCV.

Thus, novel human monoclonal antibodies to HCV E2 antigen are disclosed. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

We Claim:

1. A recombinant human monoclonal antibody that exhibits immunological binding affinity for a hepatitis C virus (HCV) E2 antigen, wherein said monoclonal antibody comprises an amino acid sequence homologous to the binding portion of a human antibody Fab molecule obtained from a combinatorial antibody library.

10

2. The monoclonal antibody of claim 1, further characterized in that said monoclonal antibody is cross-reactive to HCV E2 antigens from two or more HCV genotypes.

15

3. The monoclonal antibody of claim 2, further characterized in that said monoclonal antibody is cross-reactive to E2 antigens from HCV genotypes 1a and 2b.

20

4. The monoclonal antibody of claim 1, wherein the Fab molecule comprises a heavy chain variable region (V_H) amino acid sequence selected from the group consisting of the sequences depicted in Figure 1A (SEQ ID NO:), Figure 1B (SEQ ID NO:), Figure 1C (SEQ ID NO:), Figure 1D (SEQ ID NO:), Figure 1E (SEQ ID NO:), Figure 1F (SEQ ID NO:), and Figure 1G (SEQ ID NO:).

25

5. The monoclonal antibody of claim 1, wherein the Fab molecule comprises a light chain variable region (V_L) amino acid sequence selected from the group consisting of the sequences depicted in Figure 2A (SEQ ID NO:), Figure 2B (SEQ ID NO:), Figure 2C (SEQ ID NO:), Figure 2D (SEQ ID NO:), Figure 2E (SEQ ID NO:), Figure 2F (SEQ ID NO:), and Figure 2G (SEQ ID NO:).

30

35

6. The monoclonal antibody of claim 1, further characterized in that said monoclonal antibody has a binding affinity of at least about 1.0×10^7 .

5 7. The monoclonal antibody of claim 6, further characterized in that said monoclonal antibody also binds to the hepatitis C virus (HCV) E1/E2 complex as determined by an ELISA assay.

10 8. The monoclonal antibody of claim 1, wherein said monoclonal antibody comprises a $F(ab')_2$ fragment capable of binding to a hepatitis C virus (HCV) E2 antigen.

15 9. The monoclonal antibody of claim 1, wherein said monoclonal antibody consists essentially of a Fab molecule capable of binding to a hepatitis C virus (HCV) E2 antigen.

20 10. The monoclonal antibody of claim 9, further characterized in that said monoclonal antibody also binds to a hepatitis C virus (HCV) E1/E2 complex as determined by an ELISA assay.

25 11. A single chain Fv (sFv) molecule that exhibits immunological binding affinity for a hepatitis C virus (HCV) E2 antigen, wherein said sFv molecule comprises a binding portion formed from amino acid sequences homologous to a human antibody Fab
30 molecule obtained from a combinatorial antibody library.

12. An isolated nucleic acid molecule, comprising:
35 a first nucleotide sequence encoding a first polypeptide that is homologous to the binding portion of a $\gamma 1$ heavy chain variable region (V_H) of a human Fab

molecule that exhibits immunological binding affinity for a hepatitis C virus (HCV) E2 antigen; and

5 a second nucleotide sequence encoding a second polypeptide that is homologous to the binding portion of a κ light chain variable region (V_L) of a human Fab molecule that exhibits immunological binding affinity for a hepatitis C virus (HCV) E2 antigen.

10 13. The nucleic acid molecule of claim 12, further comprising:

15 a third nucleotide sequence encoding a first leader sequence peptide, wherein said third nucleotide sequence is operably linked to the 5' terminus of the first nucleotide sequence and is capable of causing secretion of the first polypeptide when the first polypeptide and the first leader sequence peptide are expressed; and

20 a fourth nucleotide sequence encoding a second leader sequence peptide, wherein said fourth nucleotide sequence is operably linked to the 5' terminus of the second nucleotide sequence and is capable of causing secretion of the second polypeptide when the second polypeptide and the second leader sequence peptide are expressed.

25

30 14. The nucleic acid molecule of claim 13, wherein the first nucleotide sequence is homologous to a nucleotide sequence selected from the group consisting of the sequences depicted in Figure 4A (SEQ ID NO:), Figure 4B (SEQ ID NO:), Figure 4C (SEQ ID NO:), Figure 4D (SEQ ID NO:), Figure 4E (SEQ ID NO:), Figure 4F (SEQ ID NO:), and Figure 4G (SEQ ID NO:).

35 15. The nucleic acid molecule of claim 13, wherein the second nucleotide sequence is homologous to a nucleotide sequence selected from the group

consisting of the sequences depicted in Figure 3A (SEQ
ID NO:), Figure 3B (SEQ ID NO:), Figure 3C (SEQ ID
NO:), Figure 3D (SEQ ID NO:), Figure 3E (SEQ ID
NO:), Figure 3F (SEQ ID NO:), and Figure 3G (SEQ
5 ID NO:).

16. An isolated nucleic acid molecule,
comprising a first nucleotide sequence encoding a
first polypeptide that is homologous to the binding
10 portion of a γ 1 heavy chain variable region (V_H) of a
human Fab molecule obtained from a combinatorial
library, wherein said Fab molecule exhibits
immunological binding affinity for a hepatitis C virus
(HCV) E2 antigen.

15

17. The nucleic acid molecule of claim 16,
wherein the first nucleotide sequence is homologous to
a nucleotide sequence selected from the group
consisting of the sequences depicted in Figure 4A (SEQ
20 ID NO:), Figure 4B (SEQ ID NO:), Figure 4C (SEQ ID
NO:), Figure 4D (SEQ ID NO:), Figure 4E (SEQ ID
NO:), Figure 4F (SEQ ID NO:), and Figure 4G (SEQ
ID NO:).

25

18. An isolated nucleic acid molecule,
comprising a first nucleotide sequence encoding a
first polypeptide that is homologous to the binding
portion of a κ light chain variable region (V_L) of a
human Fab molecule obtained from a combinatorial
30 library, wherein said Fab molecule exhibits
immunological binding affinity for a hepatitis C virus
(HCV) E2 antigen.

19. The nucleic acid molecule of claim 18,
35 wherein the first nucleotide sequence is homologous to
a nucleotide sequence selected from the group
consisting of the sequences depicted in Figure 3A (SEQ

ID NO:), Figure 3B (SEQ ID NO:), Figure 3C (SEQ ID NO:), Figure 3D (SEQ ID NO:), Figure 3E (SEQ ID NO:), Figure 3F (SEQ ID NO:), and Figure 3G (SEQ ID NO:).

5

20. An expression vector, comprising the nucleic acid molecule of any one of claims 12, 16 or 18 operably linked to control sequences that direct the transcription of the first and second nucleotide sequences whereby said first and second nucleotide sequences can be transcribed and translated in a host cell.

21. The expression vector of claim 20, wherein the control sequences are capable of directing the transcription of the first and second nucleotide sequences in a prokaryotic host cell.

22. The expression vector of claim 20, wherein the control sequences are capable of directing the transcription of the first and second nucleotide sequences in a eukaryotic host cell.

23. A prokaryotic host cell transformed with the expression vector of claim 21.

24. A eukaryotic host cell transformed with the expression vector of claim 22.

25. A method of producing a recombinant Fab molecule, comprising:

(a) providing a population of transformed host cells according to claim 24; and

(b) expressing said recombinant Fab molecule from the expression vector.

26. A vaccine composition comprising the

monoclonal antibody of any one of claims 1-10 and a pharmaceutically acceptable vehicle.

27. A method for providing an antibody
5 titer to HCV in a mammalian subject, comprising
introducing a therapeutically effective amount of the
vaccine composition of claim 26 to said subject.

28. A method of providing passive immunity
10 against HCV infection in a mammalian subject,
comprising introducing a therapeutically effective
amount of the vaccine composition of claim 26 to said
subject.

29. A method of treating a mammalian
15 subject having an HCV infection, comprising
introducing a therapeutically effective amount of the
vaccine composition of claim 26 to said subject.

30. The method of claim 29, further
20 comprising the step of administering a therapeutically
effective amount of α -interferon to the mammalian
subject along with said vaccine composition.

31. The method of claim 29, further
25 comprising the step of administering a therapeutically
effective amount of Ribavirin to the mammalian subject
along with said vaccine composition.

32. The method of claim 30, further
30 comprising the step of administering a therapeutically
effective amount of Ribavirin to the mammalian subject
along with said α -interferon and said vaccine
composition.

35

33. A binding complex, comprising the monoclonal antibody of any one of claims 1-10 and a detectable moiety attached thereto.

5 34. A binding complex, comprising the sFv molecule of claim 11 and a detectable moiety attached thereto.

10 35. The binding complex of claim 33 or 34, wherein the detectable moiety is selected from the group consisting of radioactive isotopes, fluorescers, chemiluminescers, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors and dyes.

15 36. A method for detecting the presence of HCV particles in a sample suspected of containing HCV, comprising:

 (a) incubating the sample with the binding complex of claim 35, wherein the incubating is
20 conducted under conditions which allow for formation of an antibody-antigen complex; and

 (b) detecting the presence or absence of the antibody-antigen complex.

25 37. A method for monitoring the progress of an anti-HCV therapeutic treatment of a HCV-infected mammalian subject, comprising:

 (a) obtaining a biological sample from said subject;

30 (b) incubating the sample with the binding complex of claim 35, wherein the incubating is conducted under conditions which allow for formation of an antibody-antigen complex; and

 (c) detecting the presence or absence of
35 the antibody-antigen complex.

1/13

FIGURE 1A (1:5, HEAVY CHAIN) [SEQ ID NO.: 1]

EVQLLEQSGA EVRKPGSSVK VSCKASGGTF SGHVTTWVRQ ^{CDR1} APCQGLENMG ESPIEGSAN YAONYAOKER ^{CDR2} DRVSIADDES TSTSFIELSN LRSDDTAVVY CARDPPRYCS AGRCYRGEEQ ^{CDR3} QWGGGLTVTVSS

EVQLLEQSGA EVKKPGSSVK VSCQVFGDTF SRITIQWLRLQ ^{CDR1} APGQGPEWNG NIIPVYNTIN YAOKEQGRLS ^{CDR2} ITADDSTSTA YNIELSSLRSE DTAVYFCARV VIPNAIRHTM ^{CDR3} GGYEDYWGQG TLVTVSS

FIGURE 1B (1:7, HEAVY CHAIN) [SEQ ID NO.: 1]

2/13

FIGURE 1C (1:11, HEAVY CHAIN) [SEQ ID NO.: 1]

EVQLLEQSGA EVKKGSSVK VSKKASGGTF SGHVSWVRQ ^{CDR1}APQGLEWVIG ^{CDR2}GSISEEGISN SAQKEQGRVS ITADESASTA YNIELSLRSE DTAIYCAKD ^{CDR3}PPRECSGGNC YPGFEQQWVQ GTILVTVSS

FIGURE 1D (L3, HEAVY CHAIN) [SEQ ID NO.: 1]

EVQLLESGG VVQGRSLRL SCAASGFTFK IYGMHWVRQ ^{CDR1}PGKGLEWVAG ^{CDR2}ISEDGSNOY ADSYKGRFV SRDNSRDTVF LQMSLRLED TAVYCATG ^{CDR3}SPEGSIKGRY YLENWVGQ TL VTVSS

3/13

FIGURE 1E (L1, HEAVY CHAIN) [SEQ ID NO.: ____]

EVQLLESGGG VVQPGRLRL SCAASGFTFS ^{CDR 1}AYGMHWVRQA ^{CDR 2}PGKGLEWVAG IWFEDGSNQYYSDSVKGRFTV .
^{CDR 3}SRDNSRNTLF LQMNSLRPED TAVYYCATEV LFESIKGRYY LENWGQGTLVTVSS

FIGURE 1F (A8, HEAVY CHAIN) [SEQ ID NO.: ____]

EVQLLESGPG LVKPSGTL^{CDR 1}SL TCTVSGGSIR SSHWWSWVRQ ^{CDR 2}PPKGLEWIG EVFESGSTIYNPSLNDRVFM .
^{CDR 3}SVDKSKDQVS LRLSSVTAAD TAVYYCARSP IKMNQGRMML DAFDIWGQGTLVIVSS

FIGURE 1G (A12, HEAVY CHAIN) [SEQ ID NO.: ____]

EVQLLESGSE VKKPGSSVKV SCRASGGGSR ^{CDR 1}SYNFENWVRQA ^{CDR 2}PGQGLEWMGG IIPMEGTANYAQKFQGRVTI .
^{CDR 3}TADESTATGY MELSSLRSED TAVYYCAMPY PKHCSHGSCW GWEDPWGQGTLVTVSS

4/13

FIGURE 2A (1:5, LIGHT CHAIN) [SEQ ID NO.:]

AELTQSPGTL SLSPGERATL SCRASOSYSS NYLAWYQQRP GQAPRLIIYG ASSRAIGIPD RESGSGSGTD FTLTISRLEP EDFAVYVCOLYGNSRWTFGGQ GTKVEIK
 CDR1 CDR 2 CDR3

AELTQSPATL SLSPGERATL SCRASOSYVK YLAWYQKPG QAPRLIIYDA SNRAIGIPAR FSGSGSGTDF TLTISNLEPE DFAVYCCQRSDWVTFGGGT KVEIK
 CDR 1 CDR 2 CDR3

FIGURE 2B (1:7, LIGHT CHAIN) [SEQ ID NO.:]

5/13

FIGURE 2C (1:11, LIGHT CHAIN) [SEQ ID NO.:]

CDR1
 AELTQSPGTL SLSPGERATL SCGASOSVRS NYLAWYQQKP GOAPRLIYG VSSRAIGIPD RESGSGSGTD FTLTISRLEP EDFAVYVCQQ YGSSPRIFGQ GTKLEIK
 CDR2
 CDR3

CDR1
 AELTQSPATL SVSPGERASL SCRASOSVGN NLAWYQQKPG QAPRLIYG NTRAIGTPDR FSGSGSGTEF TLTISSIQSE DEAVYFCQHY STWPLIFGGG TKVEFK
 CDR2
 CDR3

FIGURE 2D (L3, LIGHT CHAIN) [SEQ ID NO.:]

6/13

FIGURE 2E (L1, LIGHT CHAIN) [SEQ ID NO.:]

AELTQSPGTL SLSVGERATL SCRASQNIYS ^{CDR 1}GYLGWYQQKP ^{CDR 2}GPPRLIYG ASNRATGIPD
 - RFSGSGSGTD FTLTISRLES ^{CDR 3}EDFAVYCCQ YGSPPYTFGQ GTKVEIK

FIGURE 2F (A8, LIGHT CHAIN) [SEQ ID NO.:]

AELTQSPSSL SAFVGDRVIT ^{CDR 1}TCRASQSISR ^{CDR 2}NLNWYQQKPG TAPKVLIAA SSLQSGVPSR
 - FSGSGSGIDF ^{CDR 3}TLTITSLQPE DFATYCCQS YIIPRIFGQG TKVEIK

FIGURE 2G (A12, LIGHT CHAIN) [SEQ ID NO.:]

AELTQSPGTL SLSVGERATL SCRASQSLSS ^{CDR 1}KYLAWYQQKP ^{CDR 2}GQAPRLFIYD ASSRATGIPD
 - RFSGSGSGTD FTLTISRLEP ^{CDR 3}EDFAVYCCQ YGIPRIFGQG TKVEIK

7/13

FIGURE 3A (1:5, LIGHT CHAIN) [SEQ ID NO.:]

GAGCTCAGC AGTCTCCAGG CACCCTGTCT TTGTCTCCAG GGGAAAGAGC CACCCTCTCC TGCAGGGCCA
GTCAGAGTGT TAGCAGCAAT TACTTAGCCT GGTACCAAGCA GAGACCTGGC CAGGCTCCCA GGCTCCTCAT
CTATGGTGCA TCCAGCAGGG CCACTGGCAT CCCAGACAGG TTCAGTGCCA GTGGGTCTGG GACAGACTTC
ACTCTACCA TCAGCAGACT GGAGCCTGAA GATTTGCAG TGTATTACTG TCAGCTTTAT GGTAACCTCAC
GTTGGACGTT CGGCCAAGG ACCAAGGTGG AGATCAAA

GAGCTCACTC AGTCTCCAGC CACCCTGTCT TTGTCTCCAG GGGAAAGAGC CACCCTCTCC TGCAGGGCCA
GTCAGAGTGT TAACAAGTAC TTAGCCTGGT ACCAACAGAA ACCTGGCCAG GCTCCCAGGC TCCTCATCTA
TGATGCATCC AACAGGGCCA CTGGCATCCC AGCCAGGTTT AGTGGCAGTG GGTCTGGGAC AGACTTCACT
CTCACCATCA GCAACCTAGA GCCTGAAGAT TTGTCAGTTT ATTACTGTCA GCAGCGTAGC GACTGGGTCA
CTTTCGGCGG AGGGACCAAG GTGGAGATCA AA

FIGURE 3B (1:7, LIGHT CHAIN) [SEQ ID NO.:]

8/13

FIGURE 3C (1:11, LIGHT CHAIN) [SEQ ID NO.: 1]

GAGCTCAGC AGTCTCCAGG CACCCTGTCT TTGTCTCCAG GGAAGAGC CACCCTCTCC TCGGGGCCA
GTCAGAGTGT TAGGAGCAAC TACTTAGCCT GTTACCAGCA AAACCTGGC CAGGCTCCCA GGCTCCTCAT
CTATGGTGT TCCAGCAGG CCACTGGCAT CCGACACAGG TTCAGTGGCA GTGGTCTGG GACAGACTTC
ACTCTACCA TCAGCAGACT GGAGCCTGAA GATTTGCAG TGTATTACTG TCAGCAGTAT GGTAGCTCAC
CTCGGACTTT TGGCCAGGG ACCAAGTTGG AGATCAAA

GAGCTCAGC AGTCTCCAG CACCCTGTCT GTGTCTCCAG GGAAGAGC CTCCCTCTCC TGCAGGGCCA
GTCAGAGTGT CCGTAACAAT TTAGCTTGGT ATCAGCAGAA ACCTGGCCAG GCTCCAGGC TCCTCATTTA
TGGTGGAAAC ACCAGAGCCA CTGGTACCCC AGACAGGTTT AGTGGCAGTG GGTCTGGGAC AGAATTCAC
CTCACCATCA GCAGCCTGCA GTCTGAGGAC TTTCAGTTT ATTCTGTCA ACACTATAGT ACCTGGCCGC
TCACTTTCGG CGGGGGGACC AAGTCGAGT TCAAG

FIGURE 3D (L3, LIGHT CHAIN) [SEQ ID NO.:]

9/13

FIGURE 3E (L1, LIGHT CHAIN) [SEQ ID NO.: ____]

GAGGTGCAGC TGCTCGAGTC TGGGGAGGC GTGGTCCAGC CTGGGAGGTC CTTGAGACTC TCCTGTGCAG
 CGTCTGGATT CACCTTCAGT GCTTATGGCA TGCACCTGGT CGCCAGGCT CCAGGCAAGG GGCTGGAGTG
 GGTGGCAGGT ATATGGTTTG ATGGAAGTAA TCAATACTAT TCAGACTCCG TGAAGGGCCG ATTCACCGTC
 TCCAGAGACA ATCCAGGAA CACGCTGTT CTGCAAAATGA ACAGCCTGAG ACCCGAGGAC ACGGCTGTCT
 ATTACTGTGC GACAGAGGTA CTTTITGGAT CGATTAAGGG GCGTTACTAC CTTGAAAACCT GGGGCCAGGG
 AACCCTGGTC ACCGTCTCCT CA

FIGURE 3F (A8, LIGHT CHAIN) [SEQ ID NO.: ____]

GGGGAGCTCA CCAGTCTCC ATCGTCCCTG TCTGCATTG TNGGAGACAG AGTCACCATC ACTTGCCGGG
 CAAGTCAGAG TATTAGCAGG AACTTAAAT GGTATCAGCA GAAACCAGGG ACAGCCCTTA AGTCTCTGAT
 CTATGCTGCA TCCAGTTTGC AAAGTGGGT CCCATCGAGG TTCAGTGGCA GTGGATCTGG GACAGATTTC
 ACTCTACCA TCACCACTCT GCAACCTGAA GATTTTGCA CTTACTATTG TCAACAGAGT TACACAACCC
 CTCGGACGTT CGGCCAAGGG ACCAAGGTGG AAGTCAAA

FIGURE 3G (A12, LIGHT CHAIN) [SEQ ID NO.: ____]

GGCAGCTCA CGCAGTCTCC AGGCACCCTG TCTTGTCTC CAGGGGAAAG AGCCACCCTC TCCTGCAGGG
 CCAGTCAGAG TCTTAGCAGC AAATACTTAG CNTGGTACCA ACAGAAACCT GGCCAGGCTC CCAGGCTCTT
 CATTATGAT GCATCCAGCA GGGCCACTGG CATCCCAGAC AGGTTTCAGTG GCAGTGGGTC TGGGACAGAC
 TTCACCTCTCA GCATCAGCAG ATTGGAGCCT GAAGATTTTG CAGTGTATTA CTGTCAGCAG TATGGAACAC
 CTCGCACCTT CGGCCAGGGG ACCAAGGTGG AAATCAAA

10/13

FIGURE 4A (1:5, HEAVY CHAIN) [SEQ ID NO.:]

CTCGAGCAGT CTGGGGCTGA GGTGAGGAAG CCTGGGTCCT CGGTGAAGGT CTCCTGCAAG GCTTCTGGAG
 GCACCTTCAG CGGCCATGTT ATCACCTGGG TCGACAGGC CCTGGACAA GGACTTGAGT GGATGGGAGA
 GAGCATCCCT ATCTTTGGTT CCGCAAACTA CGCTCAAAAC TACGCTCAGA AATTCCGGGA CAGAGTCTCG
 ATTATCGCGG ACGAATCCAC GAGCACGTGG TTCAATTGAGC TGAGCAACCT GAGATCTGAC GACACGGCCG
 TCTACTACTG TCGGAGAGAC CCTCCAAGAT ATTGCACTGC TGGTAGATGC TACCCGGGAT TCTTCCAGCA
 GTGGGGCCAG GGCACCCCTCG TCACCGTCTC CTCA

CTCGAGCAGT CTGGGGCTGA GGTGAAGAAG CCTGGGTCCT CGGTGAAGGT CTCCTGTGAG GTTTTGGAG
 ACACCTTCAG CAGATACACT ATTCAGTGGT TCGACAGGC CCTGGACAA GGCCTGAGT GGATGGGAAA
 TATCATCCCT GTCTATAATA CACCAAACTA CGCGCAGAAG TTTCAGGGA GACTCTCGAT AACCGCCGAC
 GATTCCACGA GCACAGCCTA CATGGAACCTG AGTAGCCTCA GATCTGAGGA CACGGCCGTC TATTTCTGTG
 CGAGAGTCGT AATACCAAAT GCAATCCGGC ACACGATGGG ATATTACTTT GACTACTGGG GCCAGGGAAC
 CCTGGTCACC GTCTCCTCA

FIGURE 4B (1:7, HEAVY CHAIN) [SEQ ID NO.:]

11/13

FIGURE 4C (1:11, HEAVY CHAIN) [SEQ ID NO.]

CTCGAGCAGT CTGGGGCTGA GGTGAAGAAG CCTGGGTCCT CAGTGAAGT CTCCTGCAAG GCTTCTGGAG
GCACCTTCAG CGGCCATGTT ATCAGCTGGG TCGACAGGC CCTGGACAA GGCCTTGAGT GGATGGGGGG
GAGTATCTCT TTCTTTGGCA CATCAAACTC CGCACAGAAG TTCCAGGGCA GAGTCTCGAT TACCGCGGAC
GAATCCGGCA GCACAGCCTA CATGGAGCTG AGTAGCCTGA GATCGGAGGA CACGGCCATC TATTACTGTG
CGAAAGACCC TCCAAGATTT TGTAGTGGTG GTAACCTGCTA CCCGGGGTTC TTCCAGCAGT GGGGCCAGGG
CACCTTGGTC ACCGTCTCCT CA

FIGURE 4D (L3, HEAVY CHAIN) [SEQ ID NO.:]

CTCGAGTCGG GGGAGGGCGT GGTCCAGCCT GGGAGGTCCC TGAGACTCTC CTGTGCAGCG TCTGGATTCA
CCTTCAAGAC GTATGGCATG CACTGGGTCC GCCAGGCTCC AGGCAAGGG CTGGAGTGGG TGGCAGGTAT
TTCGTTTGAT GGAAGTAACC AATATTACGC AGACTCCGTG AAGGGCCGAT TCATCGTCTC CAGAGACAAT
TCCAGGGACA CGGTGTTTCT GCAGATGAGC AGCCTGAGAC TCGAGGACAC GGCTGTCTAT TACTGTGCGA
CAGAGGGTTC TCCTTTTGGC TCGATTAAAG GCGGTTACTA CCTTGAAAT TGGGGCCAGG GAACCCCTGGT
CACCGTCTCC TCA

12/13

FIGURE 4E (L1, HEAVY CHAIN) [SEQ ID NO.: ____]

GAGGTGCAGC TGCTCGAGTC TGGGGGAGGC GTGGTCCAGC CTGGGAGGTC CCTGAGACTC TCCTGTGCAG
 CGTCTGGATT CACCTTCAGT GCCTATGGCA TGCACCTGGGT CCGCCAGGCT CCAGGCAAGG GGCTGGAGTG
 GGTGGCAGGT ATATGGTTTG ATGGAAGTAA TCAATACTAT TCAGACTCCG TGAAGGGCCG ATTCACCGTC
 TCCAGAGACA ATCCAGGAA CACGCTGTTT CTGCAAAATGA ACAGCCTGAG ACCGAGGAC ACGGCTGTCT
 ATTACTGTGC GACAGAGGTA CTTTTTGGAT CGATTAGGG GCGTTACTAC CTTGAAACT GGGGCCAGGG
 AACCCCTGGTC ACCGTCTCCT CA

FIGURE 4F (A8, HEAVY CHAIN) [SEQ ID NO.: ____]


GAGGTGCAGC TGCTCGAGTC GGGCCAGGA CTGGTGAAGC CTTGGGGAC CCTGTCCCTC ACCTGCACCTG
 TCTCTGGTGG CTCCATCAGG AGCAGTCACT GGTGGAGTTG GGTCGCCAG CCCCAGGGA AGGGACTGGA
 GTGGATTGGA GAAGTCTTTT TTAGTGGAAG CACCATCTAC AACCCATCCC TCAACGATCG AGTCTTCATG
 TCTGTAGACA AGTCCAAAGGA CCAGGTCTCC CTGAGGCTGA GCTCTGTGAC CGCCGCGGAC ACGGCCGTGT
 ATTACTGTGC GAGATCCCC ATAAAATGA ATCAGGGAAG AATGATGTTG GATGCCCTTG ATATCTGGGG
 CCAGGGGACA CTCGTCATCG TCTCTTCC

FIGURE 4G (A12, HEAVY CHAIN) [SEQ ID NO.: ____]


GAGGTGCAGC TGCTCGAGTC TGGGTCTGAG GTGAAGAAGC CTGGGTCTTC GGTGAAGTC TCCTGCAGGG
 CCTCTGGAGG CAGCTTCAGA AGCTACAATT TCAATTGGGT GCGACAGGCC CCTGGACAAG GTCTTGAGTG
 GATGGGAGGC ATCATCCCTA TGTTCCGAAC AGCAAACTAC GCACAGAAGT TTCAGGGCAG AGTCACAATT
 ACCGGGACG AATCCACGGC CACAGGCTAC ATGGAGTTGA GCAGTCTGAG ATCTGAAGAC ACGGCCGTTT
 ATTACTGTGC GATGCCCTAT CCAAAACATT GCAGTCGTGG AAGTTGCTGG GGCTGGTTCC ACCCCTGGGG
 CCAGGGAAC CTGGTCACCG TGTCTTCA

13/13

% Neutralisation of E2 (HCV-1a) binding



1:5



L3

1:7

1:11

FIGURE 5

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 97/01977

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/62 C07K16/10 C12N15/13 C12N15/70 C12N15/85
 C12N1/21 C12N5/10 A61K39/42 G01N33/577 G01N33/569
 A61K47/48 A61K38/20 A61K31/70 //(A61K39/42,38:21),

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>D. ROSA ET AL.: "A quantitative test to estimate neutralizing antibodies to the hepatitis C virus: Cytofluorimetric assessment of envelope glycoprotein 2 binding to target cells."</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 93, no. 5, 5 March 1996, WASHINGTON, DC, USA, pages 1759-1763, XP000615446 cited in the application see abstract see page 1760, left-hand column, line 13 - line 28</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-25, 33-37

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

& document member of the same patent family

Date of the actual completion of the international search

17 September 1997

Date of mailing of the international search report

29.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Nooij, F

INTERNATIONAL SEARCH REPORT

International / cation No

PCT/EP 97/01977

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 (A61K39/42,31:70)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	M. PERSSON ET AL.: "Generation of diverse high-affinity human monoclonal antibodies by repertoire cloning." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 88, no. 6, 15 March 1991, WASHINGTON, DC, USA, pages 2432-2436, XP002040883 cited in the application see the whole document ---	1-25, 33-37
A	WO 93 04205 A (ABBOTT LABORATORIES) 4 March 1993 see the whole document --- -/--	1-11, 33-37

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

17 September 1997

Date of mailing of the international search report

29.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Nooij, F

INTERNATIONAL SEARCH REPORT

International Publication No.

PCT/EP 97/01977

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	A. PRINCE ET AL.: "Visualization of hepatitis C virions and putative defective interfering particles isolated from low-density lipoproteins." JOURNAL OF VIRAL HEPATITIS, vol. 3, no. 1, January 1996, pages 11-17, XP002040884 see abstract see page 14, right-hand column, line 11 - page 15, left-hand column, line 5 ---	1-11, 33-37
A	K. SIEMONEIT ET AL.: "Human monoclonal antibodies for the immunological characterization of a highly conserved protein domain of the hepatitis C virus glycoprotein E1." CLINICAL AND EXPERIMENTAL IMMUNOLOGY, vol. 101, no. 2, August 1995, OXFORD, GB, pages 278-283, XP002040885 see abstract ---	1-11, 33-37
P,X	F. HABERSETZER ET AL.: "Isolation of human monoclonal antibodies (HMabs) directed at conformational determinants of the hepatitis C virus (HCV) E2 envelope protein." HEPATOLOGY, SUPPLEMENT, vol. 24, no. 4 part 2, October 1996, BALTIMORE, MD, USA, page 381A XP002040886 see abstract 1020 ---	1-37
P,X	S-W. CHAN ET AL.: "Human recombinant antibodies specific for hepatitis C virus core and envelope E2 peptides from an immune phage display library." JOURNAL OF GENERAL VIROLOGY, vol. 77, no. 10, October 1996, LONDON, GB, pages 2531-2539, XP002040887 see the whole document -----	1-37

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP 97/01977

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

- 1 ☒ Claims Nos. 27-32
because they relate to subject matter not required to be searched by this Authority, namely.
Remark: Although claim(s) 27-32
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
- 2 ☐ Claims Nos.
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically
- 3 ☐ Claims Nos.
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows.

- 1 ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims
- 2 ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee
- 3 ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.
- 4 ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International / cation No

PCT/EP 97/01977

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9304205 A	04-03-93	US 5308750 A	03-05-94
		AU 2585092 A	16-03-93
		CA 2115923 A	04-03-93
		EP 0603307 A	29-06-94
		JP 6510192 T	17-11-94
